arXiv:0707.3024v2 [hep-ph] 5 Jun 2008

Viability of Dirac phase leptogenesis

Alexey Anisimov[a], Steve Blanchet[b] and Pasquale Di Bari[b]

[a] *Institut de Théorie des Phénomènes Physiques*
Ecole Polytechnique Fédérale de Lausanne
CH-1015 Lausanne, Switzerland
[b] *Max-Planck-Institut für Physik*
(Werner-Heisenberg-Institut)
Föhringer Ring 6, 80805 München, Germany

June 5, 2008

Abstract

We discuss the conditions for a non-vanishing Dirac phase δ and mixing angle θ_{13}, sources of CP violation in neutrino oscillations, to be uniquely responsible for the observed matter-antimatter asymmetry of the universe through leptogenesis. We show that this scenario, that we call δ-leptogenesis, is viable when the degenerate limit (DL) for the heavy right-handed (RH) neutrino spectrum is considered. We derive an interesting joint condition on $\sin\theta_{13}$ and the absolute neutrino mass scale that can be tested in future neutrino oscillation experiments. In the limit of hierarchical heavy RH neutrino spectrum (HL), we strengthen the previous result that δ-leptogenesis is only very marginally allowed, even when the production from the two heavier RH neutrinos is taken into account. An improved experimental upper bound on $\sin\theta_{13}$ and (or) an account of quantum kinetic effects could completely rule out this option in the future. Therefore, δ-leptogenesis can be also regarded as a motivation for models with degenerate heavy neutrino spectrum.

1 Introduction

Leptogenesis [1], a cosmological consequence of the see-saw mechanism [2], provides an attractive explanation for the baryon asymmetry of the Universe, one of the most long-standing cosmological puzzles. A lepton asymmetry is produced in the decays of the very heavy RH neutrinos predicted by the see-saw mechanism. In order for the ($B-L$ conserving) sphaleron processes to be able to convert part of the lepton asymmetry into a baryon asymmetry, very high temperatures, $T \gtrsim M_{\rm ew} \sim 100\,{\rm GeV}$, are required in the early Universe [3].

In comparison with other models of baryogenesis, leptogenesis offers the unique advantage of relying on an ingredient of physics beyond the Standard Model, neutrino masses, already confirmed by the experiments. Furthermore and very interestingly, a quantitative analysis [4] shows that the values of the atmospheric and of the solar neutrino mass scales, inferred from neutrino mixing experiments, favor leptogenesis to work in a mildly 'strong wash-out regime': inverse processes are strong enough to wash-out any contribution to the final asymmetry depending on the initial conditions but not too strong to prevent successful leptogenesis. In this way the observed matter-antimatter asymmetry can be unambiguously explained within a minimal extension of the SM where RH neutrinos with a Majorana mass term and Yukawa couplings are added to the SM Lagrangian and the see-saw limit is assumed. No particular assumptions on the initial conditions are required, in complete analogy with what happens in the calculation of the primordial nuclear abundances within standard Big Bang Nucleosynthesis.

In a typical N_1-dominated scenario, the asymmetry is dominantly produced from the decays of the lightest RH neutrino N_1. A necessary (but not sufficient) condition is the assumption of a mild hierarchy in the heavy neutrino mass spectrum, such that M_2, the mass of the next-to-lightest RH neutrino, is approximately three times larger than M_1, the mass of the lightest RH neutrino [5].

In an unflavored analysis, a stringent lower bound on M_1 holds [6]. In the decoupling limit, when the N_1-decay parameter $K_1 \rightarrow 0$ and assuming an initial thermal abundance, one finds [1] $M_1 \gtrsim 5\times 10^8\,{\rm GeV}$ [7]. However, there are different drawbacks for the saturation of this lower bound that is anyway strongly model dependent. A more significant and stringent lower bound, $M_1 \gtrsim 3\times 10^9\,{\rm GeV}$, is obtained at the onset of the strong wash-out regime, $K_1 \simeq K_\star \simeq 3.3$ [8, 9], where the final asymmetry does not depend on the initial conditions. This lower bound implies an associated lower bound [8, 10], $T_{\rm reh} \gtrsim 1.5\times 10^9\,{\rm GeV}$, on the value of the temperature at the beginning of the standard radiation dominated regime, the reheating temperature within inflation.

---

[1] More exactly, in [7], it was found $4\times 10^8\,{\rm GeV}$. Here we are using a slightly higher value that is obtained, as we will see, when the reduced experimental error on the baryon asymmetry and on the atmospheric neutrino mass scale is taken into account.

As we said, the assumption of a mild hierarchy in the heavy neutrino mass spectrum is not sufficient to guarantee that the N_1-dominated scenario holds. It is indeed possible that, for a proper choice of the see-saw parameters, a N_2-dominated scenario holds, where the final asymmetry is dominated by the contribution from the decays of the next-to-lightest RH neutrino N_2 [11]. In this case the lower bound on M_1 does not hold any more and is replaced by a lower bound on M_2, however still implying a lower bound on $T_{\rm reh}$.

Even when flavor effects [12, 13, 14, 15, 16, 17] are taken into account [2], these lower bounds do not get relaxed [9, 18]. In particular, flavor effects do not help to alleviate the conflict with the upper bound on the reheating temperature coming from the avoidance of the gravitino problem when a supersymmetric framework is considered [19]. On the other hand, flavor effects relax the lower bound on M_1 for $K_1 \gg K_\star$ [16] and, interestingly, it has been shown that the Dirac phase and, more efficiently, the Majorana phases can strongly enhance the amount of the relaxation [9].

Besides this effect, there is an even more interesting role played by the Majorana and Dirac phases when flavor effects are taken into account. In an unflavored analysis, the final asymmetry does not depend on the low-energy phases and this represents a limit to the possibility of further tightening the link between leptogenesis predictions and low-energy neutrino experiments.

In [15] it has been shown that, accounting for flavor effects, an asymmetry can be generated even when the total *CP* asymmetry vanishes. This is possible because flavor effects introduce an additional source of *CP* asymmetry stemming from low-energy phases. The flavor composition of the anti-lepton produced in the decay of the RH neutrino can be indeed different from the one of the *CP* conjugated lepton. In this way a new intriguing scenario arises, where the Majorana and the Dirac phases, potentially observable in low-energy neutrino experiments, could act as the unique source of *CP* violation responsible for the observed matter-antimatter asymmetry of the Universe. First calculations have been presented in [17] for particular values of M_1 and within a two RH neutrino scenario, corresponding to a specific choice of the see-saw orthogonal matrix [20].

---

[2] Flavor effects were first considered in [12], and then in [13] in the particular case of 2 RH neutrinos. However, in these papers, it was found that flavor effects can only induce small corrections to the final asymmetry compared to the unflavored case. The possibility for a large enhancement was first found in [14] in the case of resonant leptogenesis and more generally in [15, 16, 17], where the typical factor 2–3 enhancement of the final asymmetry induced by flavor effects was also first understood. As for the potential role of low energy phases in providing an additional source of *CP* violation relevant for leptogenesis, it was first discussed in [15].

In [9] it has been first shown that successful leptogenesis stemming only from low-energy phases is possible and the lower bound on M_1, with its dependence on K_1 and on the initial conditions, has been calculated for a specific choice of the see-saw orthogonal matrix in the HL. It has been found that for values of K_1 in the strong wash-out regime, the allowed region is very constrained when just Majorana phases are switched on, and it is even worse when only the Dirac phase is switched on, even for $\sin\theta_{13}$ close to its experimental upper bound. Compared to the usual cases where high-energy phases contribute to CP violation as well, the lower bounds on M_1 and on the reheating temperature get much more stringent, especially in the strong wash-out regime and in particular for values of K_1 in the range favored by neutrino mixing experiments. Therefore, one can say that the asymmetry production from low-energy phases is somehow secondary compared to the usual case when leptogenesis proceeds from the high-energy phases contained in the see-saw orthogonal matrix. This conclusion has been recently confirmed also in [21] in the context of MSSM. In [22, 23, 24], the results have been generalized for an arbitrary choice of the orthogonal matrix and of the low-energy phases but without a study of the dependence on K_1 and on the initial conditions.

In this paper we focus on the particularly interesting case of δ-leptogenesis, where the Dirac phase, which has realistic chances to be observed in neutrino mixing experiments for not too small values of $\sin\theta_{13}$, is the only non-vanishing phase. We study the dependence of the final asymmetry on the initial conditions, finding the onset of the strong wash-out regime and showing the dependence of the M_1 lower bound on the important decay parameter K_1, whose value is related to the values of the neutrino masses and at the same time determines the efficiency of the asymmetry production (involving both the production of the heavy neutrinos and the wash-out). We first obtain that, in the HL, the possibility to explain the observed asymmetry is only marginally allowed and just limited to the less relevant weak wash-out regime, when the correct condition for the validity of the fully flavored regime is taken into account [25]. Then we point out that this obstacle can be nicely circumvented going beyond the HL. Indeed, like in the unflavored case [26, 27, 5], the flavored CP asymmetries, and consequently the final $B-L$ asymmetry, get enhanced and the lower bounds on M_1 and on $T_{\rm reh}$ get relaxed. The possibility of δ-leptogenesis beyond the HL has been already studied in [23] within resonant leptogenesis [27], where the heavy neutrino mass differences are equal to the resonance widths, for initial vanishing abundance and in [28] in the context of radiative leptogenesis [29] with minimal flavor violation [30].

We perform a general analysis in the degenerate limit (DL), where at least one of the degeneracies $\delta_{ji} \equiv (M_j - M_i)/M_i \lesssim 0.01$. We show that in this case the strong wash-out regime always holds and the lower bound on M_1 can be expressed through δ_{ji} and the quantity $\Delta \equiv \sin\theta_{13}\,\sin\delta$. In the most extreme case of resonant leptogenesis this turns both into a lower bound on θ_{13} and into an upper bound on the absolute neutrino mass scale that depend on each other. In this way we find that δ-leptogenesis can indeed explain the observed matter-antimatter asymmetry of the Universe in the strong wash-out regime and therefore, like leptogenesis from high-energy phases, exhibits the same virtue of independence of the initial conditions.

In Section 2 we introduce the general framework and set the notation. In Section 3 we present the results in the HL. We confirm, in a more general way, the conclusions of [9], showing that the allowed region for δ-leptogenesis is quite restricted, especially in the strong wash-out regime and considering that the asymmetry production has to switch off for $M_1 \gtrsim 10^{12}\,\mathrm{GeV}$, when the unflavored case is recovered and CP violation from low-energy phases turns off. We also verify that this conclusion holds even when the asymmetry production from the two heavier RH neutrinos is taken into account. On the other hand, we show that a N_2-dominated scenario can also be realized in δ-leptogenesis.

We conclude that one needs to go beyond the HL for successful δ-leptogenesis in the strong wash-out regime and in any case not to be just marginally allowed. Therefore, in Section 4 we study the DL showing that successful δ-leptogenesis is possible and we find a condition that relates δ_{ji} to M_i ($j = 2, 3$ and $i = 1, 2$) and to Δ. We also find an upper bound on the absolute neutrino mass scale dependent on $\sin\theta_{13}$ that makes δ-leptogenesis falsifiable independently of the RH neutrino spectrum. In Section 5 we draw the conclusions.

2 General framework

Adding to the Standard Model three RH neutrinos with a Majorana mass term M and Yukawa couplings h, after spontaneous breaking a Dirac mass term, $m_D = v\,h$, is generated by the vev v of the Higgs boson. In the see-saw limit, $M \gg m_D$, the spectrum of neutrino mass eigenstates splits in two sets, a very heavy one, N_1, N_2 and N_3 with masses respectively $M_1 \leq M_2 \leq M_3$ and almost coinciding with the eigenvalues of M, and a light one, with masses $m_1 \leq m_2 \leq m_3$ corresponding to the eigenvalues of the light neutrino mass matrix given by the see-saw formula [2],

$$m_\nu = -m_D \frac{1}{M} m_D^T \,. \qquad (1)$$

Neutrino mixing experiments measure two light neutrino mass squared differences. In a normal scheme one has $m_3^2 - m_2^2 = \Delta m^2_{\rm atm}$ and $m_2^2 - m_1^2 = \Delta m^2_{\rm sol}$, whereas in an inverted scheme one has $m_3^2 - m_2^2 = \Delta m^2_{\rm sol}$ and $m_2^2 - m_1^2 = \Delta m^2_{\rm atm}$. For $m_1 \gg m_{\rm atm} \equiv \sqrt{\Delta m^2_{\rm atm} + \Delta m^2_{\rm sol}} = (0.052 \pm 0.002)\,{\rm eV}$ [31] the spectrum is quasi-degenerate, while for $m_1 \ll m_{\rm sol} \equiv \sqrt{\Delta m^2_{\rm sol}} = (0.0089 \pm 0.0002)\,{\rm eV}$ [31] is fully hierarchical.

In the early Universe, the decays of the heavy neutrinos into leptons and Higgs bosons produce, in general, a lepton number that is partly converted into a baryon number by sphaleron ($B - L$ conserving) processes if the temperature is higher than about 100 GeV [3].

An important role is played by the decay parameters of the heavy neutrinos defined as $K_i \equiv \widetilde{\Gamma}_i / H_{T=M_i}$, the ratios of the decay widths to the expansion rate when the RH neutrinos start to become non-relativistic at $T = M_i$. For $K_i \ll 1$ the bulk of the N_i decays occurs when they are non-relativistic and the inverse decays are not effective anymore. In this case all decays occur out-of-equilibrium and the wash-out of the asymmetry is weak. On the other hand, for $K_i \gg 1$, the heavy neutrinos decays are balanced by inverse processes. In this case the heavy neutrino abundance tracks quite closely the equilibrium abundance and the wash-out of the asymmetry is potentially, but not necessarily, strong. The answer depends on a detailed description of flavor effects that are triggered by the charged lepton Yukawa interactions with a rate $\Gamma_\alpha \simeq 5 \times 10^{-3}\, T\, f_\alpha^2\, (\alpha = e, \mu, \tau)$ [32], where the f_α's are the charged lepton Yukawa couplings in the diagonal basis.

If $\Gamma_\alpha \ll \sum_i \Gamma^i_{\rm ID}$ [3], during all the relevant period of the asymmetry generation, then the lepton state coherence is preserved between decays and inverse decays and the unflavored regime, where flavor effects are negligible, holds. This requirement implies [25]

$$M_1 \gtrsim 5 \times 10^{11}\,{\rm GeV}\,. \tag{2}$$

In the unflavored regime the condition $K_i \gg 1$ is also sufficient for the wash-out regime to be strong. It is important to stress that in this regime the only source of CP violation is due to a different total decay rate into leptons and anti-leptons and, as it is well known, it stems uniquely from high-energy phases. Therefore, in the unflavored regime, δ-leptogenesis is not viable.

If the charged lepton Yukawa interactions are in equilibrium ($\Gamma_\alpha > H$) and faster than inverse decays,

---

[3] Notice that more rigorously this condition should be written replacing the simple sum of the inverse decays rates with a sum weighted with projectors taking into account that the lepton produced by the decay of a RH neutrino N_i is different by that lepton produced by the decay of a RH neutrino $N_{j\neq i}$ and therefore is not in general fully absorbed by the $N_{j\neq i}$ inverse decay [12].

$$\Gamma_{\alpha} \gtrsim \sum_{i} \Gamma_{\rm ID}^{i} \,, \tag{3}$$

during the relevant period of the asymmetry generation, then the lepton quantum states lose coherence between the production at decay and the subsequent absorption in inverse processes. In this way the Higgs bosons interact incoherently with leptons of each flavor. In the limit case, when the quantum state becomes completely incoherent and is fully projected in one of the flavor eigenstates, each lepton flavor can be treated as a statistically independent particle species and a 'fully flavored regime' is obtained. One has to distinguish a two-flavor regime, for $M_1 \gtrsim 10^9\,{\rm GeV}$, such that the condition Eq. (3) is in any case satisfied only for $\alpha = \tau$, and a three-flavor regime, where the condition Eq. (3) applies also to $\alpha = \mu$.

In the fully (two or three) flavored regime, classic Boltzmann equations can be used like in the unflavored regime, with the difference, in general, that now each single flavor asymmetry has to be tracked independently.

In the fully flavored regime there are two new effects compared to the unflavored regime [15]. These can be understood introducing the projectors and writing them as the sum of two terms,

$$P_{i\alpha} \equiv |\langle l_i | l_\alpha \rangle|^2 = P^0_{i\alpha} + \frac{\Delta P_{i\alpha}}{2} \tag{4}$$

$$\bar{P}_{i\alpha} \equiv |\langle \bar{l}'_i | \bar{l}_\alpha \rangle|^2 = P^0_{i\alpha} - \frac{\Delta P_{i\alpha}}{2} \,. \tag{5}$$

The first effect is a reduction of the wash-out compared to the unflavored regime and is described by the tree level contribution $P^0_{i\alpha} = (P_{i\alpha} + \bar{P}_{i\alpha})/2$ setting the fraction of the total asymmetry, produced in N_i-decays, that goes into each single flavor α. In the fully flavored regime, each single inverse decay involves an independent lepton flavor eigenstate and therefore does not wash out, in general, as much asymmetry as that one produced in each single decay but an amount reduced by $P^0_{i\alpha}$.

The second effect is an additional *CP* violating contribution coming from a different flavor composition between $|l_i\rangle$ and $CP|\bar{l}'_i\rangle$. This can be described in terms of the projector differences $\Delta P_{i\alpha} \equiv P_{i\alpha} - \bar{P}_{i\alpha}$, such that $\sum_\alpha \Delta P_{i\alpha} = 0$. Indeed, defining the flavored *CP* asymmetries,

$$\varepsilon_{i\alpha} \equiv -\frac{\Gamma_{i\alpha} - \overline{\Gamma}_{i\alpha}}{\Gamma_i + \overline{\Gamma}_i} \,, \tag{6}$$

where $\Gamma_{i\alpha} \equiv P^0_{i\alpha}\,\Gamma_\alpha$ and $\bar{\Gamma}_{i\alpha} \equiv P^0_{i\alpha}\,\bar{\Gamma}_\alpha$, these can be now written as

$$\varepsilon_{i\alpha} = \varepsilon_i\, P^0_{i\alpha} + \frac{\Delta\, P_{i\alpha}}{2} \,, \tag{7}$$

where $\varepsilon_i \equiv \sum_\alpha \varepsilon_{i\alpha}$ are the total CP asymmetries. In the last expression one can see that the first term is the usual contribution due to a different decay rate into lepton and anti-leptons and the second is the additional contribution due to a possible different flavor composition between $|l_i\rangle$ and $CP|\bar{l}'_i\rangle$.

Taking into account only decays and inverse decays with proper subtraction of the resonant contribution from $\Delta L = 2$ and $\Delta L = 0$ processes [10, 14, 15], the set of effective classic Boltzmann equations valid in the fully three-flavored regime can be written as

$$\begin{aligned} \frac{dN_{N_i}}{dz} &= -D_i\,(N_{N_i} - N_{N_i}^{\rm eq}) & (i = 1, 2, 3) \\ \frac{dN_{\Delta_\alpha}}{dz} &= \sum_i \varepsilon_{i\alpha}\,D_i\,(N_{N_i} - N_{N_i}^{\rm eq}) - \sum_i P_{i\alpha}^0\,W_i^{\rm ID}\,N_{\Delta_\alpha} & (\alpha = e, \mu, \tau), \end{aligned} \tag{8}$$

where $z \equiv M_1/T$ and where we indicated with N_X any particle number or asymmetry X calculated in a portion of co-moving volume containing one heavy neutrino in ultra-relativistic thermal equilibrium, so that $N_{N_i}^{\rm eq}(T \gg M_i) = 1$. Defining $x_i \equiv M_i^2/M_1^2$ and $z_i \equiv z\,\sqrt{x_i}$, the decay factors are given by

$$D_i \equiv \frac{\Gamma_{D,i}}{H\,z} = K_i\,x_i\,z\,\left\langle\frac{1}{\gamma_i}\right\rangle\,, \tag{9}$$

where H is the expansion rate. The total decay rates, $\Gamma_{D,i} \equiv \Gamma_i + \bar{\Gamma}_i$, are the product of the decay widths times the thermally averaged dilation factors $\langle 1/\gamma_i\rangle$, given by the ratio $\mathcal{K}_1(z_i)/\mathcal{K}_2(z_i)$ of the modified Bessel functions. The equilibrium abundance and its rate are also expressed through the modified Bessel functions,

$$N_{N_i}^{\rm eq}(z_i) = \frac{1}{2}\,z_i^2\,\mathcal{K}_2(z_i)\ , \qquad \frac{dN_{N_i}^{\rm eq}}{dz_i} = -\frac{1}{2}\,z_i^2\,\mathcal{K}_1(z_i)\,. \tag{10}$$

Finally, the inverse decays wash-out terms are given by

$$W_i^{\rm ID}(z) = \frac{1}{4}\,K_i\,\sqrt{x_i}\,\mathcal{K}_1(z_i)\,z_i^3\,. \tag{11}$$

We are neglecting the non resonant contributions from $\Delta L = 2$ and $\Delta L = 0$ processes, a good approximation for $M_1 \ll 10^{14}\,{\rm GeV}\,(m_{\rm atm}^2/\sum_i m_i^2)$, as we will always consider. We are also neglecting $\Delta L = 1$ scatterings [33, 34, 35, 17], giving a correction to a level less than $\sim$ 10% [9] and spectator processes [36, 37] that, at least for a hierarchical heavy neutrino spectrum, produce a correction to a level less than $\sim$ 30% [37, 18]. In the degenerate limit it cannot be excluded that the effect of spectator processes is more relevant and further studies are required. We are also neglecting thermal corrections [10], that can give relevant (though with big theoretical uncertainties) corrections in the weak wash-out regime but negligible ones in the more important strong wash-out regime.

The evolution of the $N_{\Delta\alpha}$'s can be worked out in an integral form,

$$N_{\Delta\alpha}(z) = N_{\Delta\alpha}^{\rm in}\, e^{-\sum_i P_{i\alpha}^0 \int_{z_{\rm in}}^{z} dz'\, W_i^{\rm ID}(z')} + \sum_i \varepsilon_{i\alpha}\, \kappa_{i\alpha}(z)\,, \tag{12}$$

with the 9 efficiency factors given by

$$\kappa_{i\alpha}(z; K_i, P_{i\alpha}^0) = -\int_{z_{\rm in}}^{z} dz'\, \frac{dN_{N_i}}{dz'}\, e^{-\sum_i P_{i\alpha}^0 \int_{z'}^{z} dz''\, W_i^{\rm ID}(z'')}\,. \tag{13}$$

The total final $B-L$ asymmetry is then given by $N_{B-L}^{\rm f} = \sum_\alpha N_{\Delta_\alpha}^{\rm f}$. Finally, assuming a standard thermal history and accounting for the sphaleron converting coefficient $a_{\rm sph} \sim 1/3$, the final baryon-to-photon number ratio can be calculated as

$$\eta_B = a_{\rm sph}\, \frac{N_{B-L}^{\rm f}}{N_\gamma^{\rm rec}} \simeq 0.96 \times 10^{-2}\, N_{B-L}^{\rm f}\,, \tag{14}$$

to be compared with the measured value [38]

$$\eta_B^{\rm CMB} = (6.1 \pm 0.2) \times 10^{-10}\,. \tag{15}$$

Notice that the efficiency factors depend only on the $P_{i\alpha}^0$ but not on the differences $\Delta P_{i\alpha}$. Notice also that, in the two-flavor regime, the individual electron and muon asymmetries are replaced by one kinetic equation for the sum, $N_{\Delta_{e\mu}} \equiv N_{\Delta_\mu} + N_{\Delta_e}$, where the individual flavored CP asymmetries and projectors have also to be replaced by the their sum, namely $\varepsilon_{1\,e+\mu} \equiv \varepsilon_{1\mu} + \varepsilon_{1e}$ and $P_{1\,e+\mu}^0 \equiv P_{1\mu}^0 + P_{1e}^0$ [17]. The calculation is therefore somehow intermediate between the one-flavor approximation and the three-flavor regime, though the results are very similar to the three-flavor regime [9].

The flavored CP asymmetries are given by the following expression [26]

$$\varepsilon_{i\alpha} = \frac{3}{16\pi (h^\dagger h)_{ii}} \sum_{j\neq i} \left\{ {\rm Im}\left[h_{\alpha i}^\star h_{\alpha j} (h^\dagger h)_{ij}\right] \frac{\xi(x_j/x_i)}{\sqrt{x_j/x_i}} + \frac{2}{3(x_j/x_i - 1)} {\rm Im}\left[h_{\alpha i}^\star h_{\alpha j} (h^\dagger h)_{ji}\right] \right\}\,, \tag{16}$$

where

$$\xi(x) = \frac{2}{3}\, x \left[(1+x)\, \ln\left(\frac{1+x}{x}\right) - \frac{2-x}{1-x}\right]\,. \tag{17}$$

A parametrization of the Dirac mass matrix, particularly fruitful within leptogenesis, is obtained in terms of the see-saw orthogonal matrix Ω [20]

$$m_D = U\, D_m^{1/2}\, \Omega\, D_M^{1/2}\,, \tag{18}$$

where we defined $D_m \equiv \mathrm{diag}(m_1, m_2, m_3)$ and $D_M \equiv \mathrm{diag}(M_1, M_2, M_3)$. The matrix U diagonalizes the light neutrino mass matrix m_ν, such that $U^\dagger\, m_\nu\, U^\star = -D_m$, and it can be identified with the lepton mixing matrix in a basis where the charged lepton mass matrix is diagonal. Moreover, neglecting the effect of the running of neutrino parameters from high energy to low energy [39], one can assume that the U matrix can be identified with the PMNS matrix, partially measured in neutrino mixing experiments. For normal hierarchy we adopt the parametrization [40]

$$U = \begin{pmatrix} c_{12}\, c_{13} & s_{12}\, c_{13} & s_{13}\, e^{-i\,\delta} \\ -s_{12}\, c_{23} - c_{12}\, s_{23}\, s_{13}\, e^{i\,\delta} & c_{12}\, c_{23} - s_{12}\, s_{23}\, s_{13}\, e^{i\,\delta} & s_{23}\, c_{13} \\ s_{12}\, s_{23} - c_{12}\, c_{23}\, s_{13}\, e^{i\,\delta} & -c_{12}\, s_{23} - s_{12}\, c_{23}\, s_{13}\, e^{i\,\delta} & c_{23}\, c_{13} \end{pmatrix} \times \mathrm{diag}(\mathrm{e}^{\mathrm{i}\,\frac{\Phi_1}{2}}, \mathrm{e}^{\mathrm{i}\,\frac{\Phi_2}{2}}, 1)\,, \tag{19}$$

where $s_{ij} \equiv \sin\theta_{ij}$, $c_{ij} \equiv \cos\theta_{ij}$ and, neglecting the statistical errors, we will use $\theta_{12} = \pi/6$ and $\theta_{23} = \pi/4$, compatible with the results from neutrino mixing experiments. Moreover, we will adopt the 3σ range $s_{13} = 0 - 0.20$, allowed from a global 3ν analysis for unitary U [31], an approximation that holds with great precision in the see-saw limit with $M_i \gg 100\,\mathrm{GeV}$. Within the convention we are using, $m_1 \leq m_2 \leq m_3$, the case of inverted hierarchy corresponds is obtained performing a cyclic permutation of columns in the PMNS matrix parametrization Eq. (19), such that the i-th column becomes the $(i+1)$-th. Since we are interested in understanding whether a non-vanishing Dirac phase can be the only source of *CP* violation for successful leptogenesis, we will set the Majorana phases to zero. We will comment later on the effects of turning on the Majorana phases.

It will also prove useful to introduce the following parametrization for the see-saw orthogonal matrix,

$$\Omega(\omega_{21}, \omega_{31}, \omega_{32}) = R_{12}(\omega_{21})\ R_{13}(\omega_{31})\ R_{23}(\omega_{32})\ , \tag{20}$$

where

$$R_{12} = \begin{pmatrix} \sqrt{1-\omega_{21}^2} & -\omega_{21} & 0 \\ \omega_{21} & \sqrt{1-\omega_{21}^2} & 0 \\ 0 & 0 & 1 \end{pmatrix}\ ,\ R_{13} = \begin{pmatrix} \sqrt{1-\omega_{31}^2} & 0 & -\omega_{31} \\ 0 & 1 & 0 \\ \omega_{31} & 0 & \sqrt{1-\omega_{31}^2} \end{pmatrix}\ ,\ R_{23} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \sqrt{1-\omega_{32}^2} & -\omega_{32} \\ 0 & \omega_{32} & \sqrt{1-\omega_{32}^2} \end{pmatrix}\ . \tag{21}$$

Notice that, using the orthogonal parametrization, the decay parameters K_i can be expressed as linear combinations of the neutrino masses [41, 8]

$$K_i = \frac{\widetilde{m}_i}{m_\star} = \sum_j \frac{m_j}{m_\star}\, |\Omega_{ji}^2|\,, \tag{22}$$

where $\widetilde{m}_i \equiv (m_D^\dagger\, m_D)_{ii}/M_i$ are the effective neutrino masses [34] and where $m_\star$ is the equilibrium neutrino mass [8] given by

$$m_\star = \frac{16\,\pi^{5/2}\,\sqrt{g_*}}{3\,\sqrt{5}}\,\frac{v^2}{M_{Pl}} \simeq 1.08\times 10^{-3}\,\mathrm{eV}\,. \tag{23}$$

Barring huge phase cancellations and special forms for Ω, typically the K_i's span within the range $[K_{\rm sol}, K_{\rm atm}]$ where $K_{\rm sol}\equiv m_{\rm sol}/m_\star = 8.2\pm 0.2$ and $K_{\rm atm}\equiv m_{\rm atm}/m_\star = 48\pm 2$.

Before entering into a detailed analysis focusing on δ-leptogenesis, we want to discuss some general features concerning the fully flavored regime and in particular the possibility to have important deviations from the unflavored case. For definiteness and simplicity, we refer to the two-flavor case within the N_1-dominated scenario, so that $N_{B-L}^{\rm f}\simeq \varepsilon_{1\tau}\,\kappa_{1\tau}^{\rm f} + \varepsilon_{1,e+\mu}\,\kappa_{1,e+\mu}^{\rm f}$.

Consider first the 'democratic case', where $\Delta P_{1\alpha}=0$ and $P_{1\tau}=P_{1\,e+\mu}=1/2$. Summing the two equations for $\alpha=\tau$ and $\alpha=e+\mu$ one obtains a closed equation for the total asymmetry where the only effect compared to the unflavored case is that the wash-out is reduced by a factor two and the final asymmetry is obtained by replacing $K_1\rightarrow K_1/2$. Therefore, in the strong wash-out regime ($K_1\gg 1$), since $\kappa_{1\alpha}^{\rm f}\propto K_1^{-1.2}$ [42], one has approximately a factor two enhancement. Let us now consider $P_{1\mu}^0 < P_{1\tau}^0$, still with $\Delta P_{1\alpha}=0$. Since approximately $\kappa^{\rm f}_{1\alpha}\propto (P^0_{1\alpha})^{-1.2}$ and at the same time $\varepsilon_{1\alpha}\propto P^0_{1\alpha}$, one has that the final asymmetry stays approximately constant compared to the democratic case with the two contributions from the μ and τ flavors comparable with each other. Therefore, for vanishing $\Delta P_{1\alpha}$, flavor effects produce just $\mathcal{O}(1)$ corrections compared to the unflavored approximation.

This conclusion changes when non-vanishing $\Delta P_{1\alpha}$ are considered. In this case there are two remarkable possibilities.

The *first possibility* is the so called one-flavor dominated scenario, relying on the fact that, for $P^0_{1\alpha}\rightarrow 0$, one has $\max(\Delta P_{1\alpha})\propto\sqrt{P^0_{1\alpha}}$ [17]. Therefore, considering now for example $P_{1\tau}\ll P_{1e+\mu}\simeq 1$, one has that the asymmetry in the tauon flavor $\varepsilon_{1\tau}\,\kappa_{1\tau}\propto (P^0_{1\tau})^{-0.7}$, showing that there can be a large enhancement compared to the unflavored case in the strong wash-out regime. This brings to a strong relaxation of the lower bounds on M_1 and $T_{\rm reh}$ at $K_1\gg 1$, though, as we already said, not to a relaxation of the usual lowest bounds at $K_1\rightarrow 0$ or at $K_1\simeq K_\star$. It should also be said that, as shown in [25], the applicability of the one-flavor dominated scenario is strongly limited by the condition of validity of the fully flavored regime Eq. (3).

The *second possibility* relies on the observation that, contrarily to ε_1, the $\Delta P_{1\alpha}$'s depend on the low-energy phases as well and, even though $\varepsilon_1=0$, they do not vanish if the Dirac or the Majorana phases do not vanish. Therefore, one can have a final asymmetry originating just from low-energy phases [15]. This scenario represents, potentially, the most important novelty introduced by flavor effects compared to the unflavored picture and in the following Sections we will study it in detail, focusing on the case of δ-leptogenesis, when only the Dirac phase is switched on while Ω is real and the two Majorana phases vanish.

Before concluding this Section, we want to notice that one can have $\varepsilon_i = 0$ not only when the see-saw orthogonal matrix is real, but also when the absolute neutrino mass scale increases [16]. In this way, the low-energy phases can play an important role in circumventing the upper bound on the neutrino masses holding in the unflavored regime [43, 4]. It is however still to be assessed whether the fully flavored regime can offer a sufficient description to solve this issue. In [16] the bound was found to be completely nullified by flavor effects. In [25] it has been pointed out how this conclusion relies on a extension of the fully flavored regime beyond the regime of its validity given by the Eq. (3). In [44] the authors find that in the fully flavored regime, thanks to spectator processes, the bound holding in the unflavored regime, even though not nullified, is anyway relaxed to $m_1 \lesssim 2\,\mathrm{eV}$.

3 The hierarchical limit

Let us consider first δ-leptogenesis in the HL, such that $M_3 \gtrsim 3\,M_2 \gtrsim 3\,M_1$ [5]. In the unflavored regime, this assumption typically implies a N_1-dominated scenario, where the final asymmetry is dominated by the contribution from the lightest RH neutrino decays,

$$N_{B-L}^{\mathrm{f}} \simeq N_{B-L}^{\mathrm{f}}\big|_{N_1} \equiv \sum_{\alpha} \varepsilon_{1\alpha}\,\kappa_{1\alpha}\,. \tag{24}$$

Indeed, in general, in the HL one has two effects. The first effect is that the asymmetry production from the two heavier RH neutrinos, N_2 and N_3, is typically later on washed out by the N_1 inverse processes and $\kappa_3^{\mathrm{f}}, \kappa_2^{\mathrm{f}} \ll \kappa_1^{\mathrm{f}}$. The second effect is a consequence of the fact that the total *CP* asymmetries vanish in the limit when all particles running in the loops become massless and this yields typically $|\varepsilon_3| \ll |\varepsilon_2| \ll |\varepsilon_1|$.

However, for a particular choice of the see-saw parameters, $\Omega \simeq R_{23}$ and $m_1 \lesssim m_\star$, the contribution to the final asymmetry from the next-to-lightest RH neutrino N_2 is not only non-negligible but even dominant, giving rise to a N_2-dominated scenario [11]. Indeed for $\Omega \simeq R_{23}$ different things happen simultaneously. First, N_2, even though decoupled from N_1, is still coupled to N_3 and in the HL the total *CP* asymmetry ε_2 does not vanish, since it receives a non suppressed contribution from graphs where N_3 runs in the loops. On the other hand, now one has $\varepsilon_1 = 0$, since N_1 is essentially decoupled from the other two RH neutrinos. At the same time one also has $K_1 \ll 1$, so that the wash-out from N_1 inverse processes is negligible. The final result is that $|\varepsilon_2\,\kappa_2| \gg |\varepsilon_{i\neq 2}\,\kappa^{\rm f}_{i\neq 2}|$ and the final asymmetry is dominantly produced from N_2-decays.

Therefore, in the unflavored approximation and in the HL, a condition $w_{32} \simeq 1$ in the Ω-matrix parametrization (cf. Eq. (20)) is sufficient to have a negligible asymmetry production from the two heavier RH neutrinos and to guarantee that the N_1-dominated scenario holds. This condition is even not necessary for $m_1 \gg m_\star$, since in this case, due to the fact that $\widetilde{m}_1 \geq m_1$, one has necessarily $K_1 \gg 1$ and a wash-out from N_1-inverse processes is anyway strong enough to suppress a possible contribution to the final asymmetry produced from N_2-decays.

When flavor effects are taken into account, the domain of applicability of the N_1-dominated scenario reduces somehow. There are two aspects to be considered.

The first aspect is that the wash-out from N_1 inverse processes becomes less efficient. Indeed the projectors $P_{1\alpha}$ can considerably reduce the wash-out of the asymmetry produced in the flavor α from N_2-decays [45]. This turns the condition $m_1 \gg m_\star$ into a looser condition $m_1 \gg m_\star/P_{1\alpha}$. Another effect is that N_1 inverse processes can make part of the asymmetry produced in N_2 decays somehow orthogonal to the the wash-out from N_1 inverse processes [12, 46]. Recently, it has been also pointed out that spectator processes can lead to a reduction of the wash-out from N_1-inverse processes as well [47]. In this way the assumption $\kappa_{2\alpha} \ll \kappa_{1\alpha}$ is not valid in general.

The second aspect concerns the flavored *CP* asymmetries. In the HL, from the general expression Eq. (16), one has

$$\varepsilon_{1\alpha} \simeq \frac{3}{16\pi (h^\dagger h)_{11}} \sum_{j\neq 1} \frac{M_1}{M_j} {\rm Im}\left[h^\star_{\alpha 1} h_{\alpha j} (h^\dagger h)_{1j}\right] , \tag{25}$$

$$\varepsilon_{2\alpha} \simeq \frac{3}{16\pi (h^\dagger h)_{22}} \left\{ \frac{M_2}{M_3} {\rm Im}\left[h^\star_{\alpha 2} h_{\alpha 3} (h^\dagger h)_{23}\right] - \frac{2}{3} {\rm Im}\left[h^\star_{\alpha 2} h_{\alpha 1} (h^\dagger h)_{12}\right] \right\} , \tag{26}$$

$$\varepsilon_{3\alpha} \simeq -\frac{1}{8\,\pi (h^\dagger h)_{33}} \sum_{j\neq 3} \left\{ {\rm Im}\left[h^\star_{\alpha 3} h_{\alpha j} (h^\dagger h)_{j3}\right] \right\} . \tag{27}$$

Different comments are in order. The $\varepsilon_{1\alpha}$'s, like ε_1, vanish for $\Omega = R_{23}$ while the $\varepsilon_{2\alpha}$'s, like ε_2, do not. On the other hand, in the HL, the $\varepsilon_{2\alpha}$'s, contrarily to ε_2, are not suppressed when $\omega_{32} = 0$ (a particular example is given by $\Omega = R_{12}$) but, like ε_2, they vanish for $\Omega = R_{13}$.

This observation [9] can also potentially contribute to enlarge the domain of applicability of the N_2-dominated scenario when flavor effects are taken into account. Another interesting observation is that the $\varepsilon_{3\alpha}$'s, contrarily to ε_3, do not vanish in the HL. This could open the door even to a N_3-dominated scenario, though this is possible only for $M_3 \lesssim 10^{12}\,\mathrm{GeV}$, when flavor effects are effective in N_3 decays.

Therefore, when flavor effects are taken into account, the conditions of applicability of the N_1-dominated scenario become potentially more restrictive than in the unflavored case. There is a clear choice of the parameters, for $\Omega = R_{13}$ and $M_3 \gtrsim 10^{12}\,\mathrm{GeV}$, where the N_1-dominated scenario holds. Indeed in this case, in the HL, one has that $\varepsilon_{2\alpha}$ and ε_3 are suppressed. This can be considered somehow opposite to the case $\Omega = R_{23}$, where the N_2-dominated scenario holds [11].

In general, one can say that the asymmetry produced from the two heavier RH neutrinos is non-negligible if two conditions are satisfied. (i) The asymmetry generated from $N_{2,3}$-decays at $T \sim M_{2,3}$ has to be non-negligible compared to the asymmetry generated at $T \sim M_1$ from N_1-decays. This depends on an evaluation of the CP asymmetries $\varepsilon^{\alpha}_{2,3}$ and of the wash-out due to the same $N_{2,3}$-inverse processes. (ii) The asymmetry produced from $N_{2,3}$-decays has not to be afterwards washed-out by N_1-inverse processes. Notice that this second condition is subordinate to the first condition.

In the particular case of δ-leptogenesis, one has $\varepsilon_2 = \varepsilon_3 = 0$. This means that the first condition can be satisfied only if $M_2, M_3 \lesssim 10^{12}\,\mathrm{GeV}$ and this constitutes already an important limitation. In the following, we will consider different particular cases, verifying whether the production from the two heavier RH neutrinos can be neglected or not. We will find that the situation is actually similar to what happens in the unflavored case where, except for the case $\Omega \sim R_{23}$, a N_1-dominated scenario holds.

Let us therefore start showing in detail how to calculate the contribution to the final asymmetry from N_1-decays. The expression Eq. (13) for the $\kappa_{1\alpha}$'s can be specialized as

$$\kappa_{1\alpha}(z; K_1, P^0_{1\alpha}) = -\int_{z_{\rm in}}^{z} dz'\, \frac{dN_{N_1}}{dz'}\, e^{-P^0_{1\alpha}\int_{z'}^{z} dz''\, W_1^{\rm ID}(z'')} \,. \tag{28}$$

From the Eq. (13), extending an analytic procedure derived within the one-flavor approximation [8], one can obtain simple analytic expressions for the $\kappa^{\rm f}_{1\alpha}$'s. In the case of an initial thermal abundance ($N^{\rm in}_{N_1} = 1$), defining $K_{1\alpha} \equiv P^0_{1\alpha}\, K_1$, one has

$$\kappa^{\rm f}_{1\alpha} \simeq \kappa(K_{1\alpha}) \equiv \frac{2}{K_{1\alpha}\, z_B(K_{1\alpha})}\, \left(1 - e^{-\frac{K_{1\alpha}\, z_B(K_{1\alpha})}{2}}\right) \,, \tag{29}$$

where

$$z_B(K_{1\alpha}) \simeq 2 + 4\, K_{1\alpha}^{0.13}\, e^{-\frac{2.5}{K_{1\alpha}}} \,. \tag{30}$$

In the case of initial vanishing abundance ($N^{\rm in}_{N_1} = 0$) one has to take into account two different contributions, a negative and a positive one, so that

$$\kappa^{\rm f}_{1\alpha} = \kappa^{\rm f}_{-}(K_1, P^0_{1\alpha}) + \kappa^{\rm f}_{+}(K_1, P^0_{1\alpha}) \,, \tag{31}$$

whose analytic expressions, used to obtain all presented results, can be found in [9].

The condition for the validity of the fully flavored regime Eq. (3) can be specialized and re-cast like

$$M_1 \lesssim \frac{10^{12}\,\mathrm{GeV}}{2\,W_1^{\mathrm{ID}}(z_B(K_{1\alpha}))}. \tag{32}$$

This condition neglects the effect of $\Delta L = 1$ scatterings and of coherent scatterings, the first contributing with inverse decays to preserve the quantum state coherence, the second, conversely, in projecting it on the flavor basis [25]. Both of them can be as large as the effect from inverse decays. Moreover, in a rigorous quantum kinetic description, it is likely that other subtle effects contribute to the determination of the exact value of M_1 below which the fully flavored regime can be assumed. Therefore, the condition (32) should be regarded as a very qualitative one. In the plots showing the M_1 lower bound, we will then distinguish four regions. All plots will be cut at $M_1 = 10^{12}\,\mathrm{GeV}$, since above this value, according to the condition (2), the unflavored regime is recovered and the asymmetry production has to switch off. On the other hand, when the condition Eq. (32) is satisfied, one can expect the fully flavored regime to hold. There is an intermediate regime where a transition between the fully flavored regime and the unflavored regime takes place. This regime will be indicated in all plots with a squared region. This signals that, even though we still show the results obtained in the fully flavored regime, important corrections are expected, especially when M_1 gets close to $\sim 10^{12}\,\mathrm{GeV}$. Since this region describes a transition toward the unflavored regime, where the asymmetry production has to switch off, these corrections are expected to reduce the final asymmetry, making more stringent the lower bounds shown in the plots. Furthermore, since within current calculation, large corrections to the condition Eq. (32) cannot be excluded, we will also indicate, with a hatched region, that area where the condition Eq. (32) holds but a very conservative condition,

$$M_1 \lesssim \frac{10^{11}\,\mathrm{GeV}}{W_1^{\mathrm{ID}}(z_B(K_{1\alpha}))} \tag{33}$$

does not. In this region some corrections to the presented results cannot be excluded but the fully flavored regime should represent a good approximation.

We anticipate that, in the N_1-dominated scenario, successful leptogenesis always requires $M_1 \gtrsim 10^9\,\mathrm{GeV}$, where the two-flavor regime holds. Therefore, considering that we are assuming $\varepsilon_1 = \varepsilon_{1\tau} + \varepsilon_{1,e+\mu} = 0$, the Eq. (24) can be specialized into

$$N_{B-L}^{\mathrm{f}}\big|_{N_1} \simeq (\kappa_{1\,\tau}^{\mathrm{f}} - \kappa_{1,e+\mu}^{\mathrm{f}})\,\varepsilon_{1\tau}\,, \tag{34}$$

showing that, in order to have a non-vanishing final asymmetry it has to be $P_{1\tau}^0 \neq P_{1,e+\mu}^0$. The tree-level projectors can be expressed, through the orthogonal parametrization Eq.

(18), like

$$P^0_{1\alpha} = \frac{|\sum_j \sqrt{m_j}\, U_{\alpha j}\, \Omega_{j1}|^2}{\sum_j m_j\, |\Omega^2_{j1}|}\,, \tag{35}$$

that, from the Eq. (22), also implies

$$K_{1\alpha} = \left|\sum_j \sqrt{\frac{m_j}{m_\star}}\, U_{\alpha j}\, \Omega_{j1}\right|^2 . \tag{36}$$

Let us now calculate the flavored *CP* asymmetry $\varepsilon_{1\tau}$ from the general expression Eq. (25). In terms of the orthogonal parametrization Eq. (18), this can be re-cast as [17]

$$r_{1\alpha} = -\sum_{h,l} \frac{m_l\, \sqrt{m_l\, m_h}}{\widetilde{m}_1\, m_{\rm atm}}\, {\rm Im}[U_{\alpha h}\, U^\star_{\alpha l}\, \Omega_{h1}\, \Omega_{l1}]\,, \tag{37}$$

where we defined $r_{i\alpha} \equiv \varepsilon_{i\alpha}/\overline{\varepsilon}(M_i)$, with

$$\overline{\varepsilon}(M_i) \equiv \frac{3}{16\pi}\, \frac{M_i\, m_{\rm atm}}{v^2}\,. \tag{38}$$

For real Ω, the Eq. (37) gets specialized into [17]

$$r_{1\alpha} = -\sum_{h<l} \frac{\sqrt{m_l\, m_h}\, (m_l - m_h)}{\widetilde{m}_1\, m_{\rm atm}}\, \Omega_{h1}\, \Omega_{l1}\, {\rm Im}[U_{\alpha h}\, U^\star_{\alpha l}]\,. \tag{39}$$

Taking $\alpha = \tau$ and specifying the matrix elements $U_{\alpha j}$, from the Eq. (19), one has

$$r_{1\tau} = -\frac{m_{\rm atm}}{\widetilde{m}_1}\, [A_{12} + A_{13} + A_{23}], \tag{40}$$

where

$$\begin{aligned}
A_{12} &= -\frac{\sqrt{m_1\, m_2}\, (m_2 - m_1)}{m^2_{\rm atm}}\, \Omega_{11}\, \Omega_{21}\, {\rm Im}[(s_{12}\, s_{23} - c_{12}\, c_{23}\, s_{13}\, e^{i\,\delta}) \\
&\qquad \times (c_{12}\, s_{23} + s_{12}\, c_{23}\, s_{13}\, e^{-i\,\delta})\, e^{-\frac{i}{2}\,(\Phi_2 - \Phi_1)}]\,, \\
A_{13} &= \frac{\sqrt{m_1\, m_3}\, (m_3 - m_1)}{m^2_{\rm atm}}\, \Omega_{11}\, \Omega_{31}\, c_{23}\, c_{13}\, {\rm Im}[(s_{12}\, s_{23} - c_{12}\, c_{23}\, s_{13}\, e^{i\,\delta}) e^{\frac{i}{2}\,\Phi_1}]\,, \\
A_{23} &= -\frac{\sqrt{m_2\, m_3}\, (m_3 - m_2)}{m^2_{\rm atm}}\, \Omega_{21}\, \Omega_{31}\, c_{23}\, c_{13}\, {\rm Im}[(c_{12}\, s_{23} + s_{12}\, c_{23}\, s_{13}\, e^{i\,\delta})\, e^{\frac{i}{2}\,\Phi_2}]\,.
\end{aligned}$$

In the case of δ-leptogenesis ($\Phi_1 = \Phi_2 = 0$) these expressions further specialize into

$$\begin{aligned}
A_{12} &= \frac{\sqrt{m_1\, m_2}\, (m_2 - m_1)}{m^2_{\rm atm}}\, \Omega_{11}\, \Omega_{21}\, s_{23}\, c_{23}\, \Delta \\
A_{13} &= -\frac{\sqrt{m_1\, m_3}\, (m_3 - m_1)}{m^2_{\rm atm}}\, \Omega_{11}\, \Omega_{31}\, c^2_{23}\, c_{12}\, c_{13}\, \Delta\,, \\
A_{23} &= -\frac{\sqrt{m_2\, m_3}\, (m_3 - m_2)}{m^2_{\rm atm}}\, \Omega_{21}\, \Omega_{31}\, c^2_{23}\, s_{12}\, c_{13}\, \Delta\,,
\end{aligned}$$

where remember that $\Delta \equiv \sin\theta_{13}\,\sin\delta$.

It is now instructive to make some general considerations. Looking at the expression Eq. (34), one can see that, in order for the final $B-L$ asymmetry not to vanish, two conditions have to be simultaneously satisfied : $\varepsilon_{1\tau} \neq 0$ and $\kappa^{\rm f}_{1\tau} \neq \kappa^{\rm f}_{1,e+\mu}$. These two conditions are a specialization of the Sakharov necessary conditions to the case of δ-leptogenesis. Indeed, the first is the condition to have *CP* violation and, as one could expect, from the expressions found for the terms A_{ij}, one can have $\varepsilon_{1\tau} \neq 0$ only if $\Delta \neq 0$. The second condition is a specialization of the condition of departure from thermal equilibrium in quite a non-trivial way. Indeed, in the case of δ leptogenesis, in a full out-of-equilibrium situation where only decays are active, no final asymmetry is generated since $\varepsilon_1 = 0$, implying that there is an equal number of decays into lepton and anti leptons. However, the presence of inverse processes can remove this balance, yielding a different wash-out rate for the τ asymmetry and for the $e+\mu$ asymmetry, such that, if $K_{1\tau} \neq K_{1,e+\mu}$, one has a net lepton number dynamical generation. From the expression (36), one can see that this is possible independently of the value of the Dirac phase that, therefore, is directly responsible only for *CP* violation and not for lepton number violation, exactly as in neutrino mixing, where indeed lepton number is conserved. It should also be noticed that the $\varepsilon_{1\alpha}$'s are expressed through quantities ${\rm Im}[U_{\alpha h}\,U^{\star}_{\alpha l}]$, that are invariant under change of the PMNS matrix parametrization [48, 23]. Therefore, the final asymmetry depends correctly only on physical quantities.

Maximizing the asymmetry over all involved parameters for fixed M_1 and K_1 and imposing $\eta_B^{\rm max} \geq \eta_B^{\rm CMB}$ (cf. (14) and (15)), a lower bound on M_1 is obtained [9]

$$M_1 \geq M_1^{\rm min}(K_1) \equiv \frac{\overline{M}_1}{\kappa^{\rm f}_1(K_1)\,\xi_1^{\rm max}}\,, \tag{41}$$

where we introduced the quantity [4]

$$\overline{M}_1 \equiv \frac{16\,\pi}{3}\,\frac{N_\gamma^{\rm rec}\,v^2}{a_{\rm sph}}\,\frac{\eta_B^{\rm CMB}}{m_{\rm atm}} = (6.25 \pm 0.4)\times 10^8\,{\rm GeV} \gtrsim 5\times 10^8{\rm GeV}\,. \tag{42}$$

The last inequality gives the 3σ value that we used to obtain all the results shown in the figures. We also defined [9]

$$\xi_1 \equiv \sum_{\alpha=\tau,e+\mu} \xi_{1\alpha}\,, \qquad \text{with} \qquad \xi_{1\alpha} \equiv \frac{r_{1\alpha}\,\kappa^{\rm f}_{1\alpha}(K_{1\alpha})}{\kappa^{\rm f}_1(K_1)}\,, \tag{43}$$

---

[4] Notice that $\overline{M_1}$ gives the lower bound on M_1 in the unflavored case for initial thermal abundance and in the limit $K_1 \to 0$. Because of the improved determination of $\eta_B^{\rm CMB}/m_{\rm atm}$ from the 3 years WMAP data [38] and from new data from neutrino oscillation experiments, in particular from the MINOS experiment, the error on $\overline{M}_1$ is halved compared to the previous estimation in [8].

that gives the deviation introduced by flavor effects compared to the unflavored approximation in the hierarchical light neutrino case ($m_1 = 0$). Notice that $r_{1\tau} \propto \Delta$, implying $N^{\rm f}_{B-L} \propto \Delta$ as well. Therefore, the maximum asymmetry is obtained for $|\delta| = \pi/2$ and $s_{13} = 0.20$.

The calculation of the contribution to the asymmetry from N_2-decays proceeds in an analogous way. Again this can always be calculated in the two-flavor regime, since, in the HL, successful leptogenesis always implies $M_2 \gtrsim 10^9\,{\rm GeV}$. Therefore, one can write an expression similar to the Eq. (34) for the contribution to the final asymmetry from N_2-decays,

$$N^{\rm f}_{B-L}\big|_{N_2} \simeq (\kappa^{\rm f}_{2\,\tau} - \kappa^{\rm f}_{2,e+\mu})\,\varepsilon_{2\tau}\,. \tag{44}$$

The difference is now in the calculation of the efficiency factors that are suppressed by the wash-out of the N_1 inverse processes. In the HL this additional wash-out factorizes and [5, 9, 45]

$$\kappa^{\rm f}_{2\alpha} \simeq \kappa(K_{2\alpha})\, e^{-\frac{3\pi}{8}\,K_{1\alpha}}\,, \tag{45}$$

where $K_{2\alpha} \equiv P^0_{2\alpha}\,K_2$. For the calculation of the tree-level projectors $P^0_{2\alpha}$ an expression analogous to the Eq. (35) holds.

The calculation of the contribution to the final asymmetry from N_3-decays proceeds in a similar way and analogous expressions hold. The only non trivial difference is that now, in the calculation of the efficiency factors, one has also to include the wash-out from the N_2 inverse processes, so that

$$\kappa^{\rm f}_{3\alpha} \simeq \kappa(K_{3\alpha})\, e^{-\frac{3\pi}{8}\,(K_{1\alpha}+K_{2\alpha})}\,. \tag{46}$$

Notice that in the calculation of $\kappa^{\rm f}_{2\alpha}$ ($\kappa^{\rm f}_{3\alpha}$) we are not including a possible effect where part of the asymmetry in the flavor $\alpha = e+\mu$ produced in N_2 (N_3-decays) is orthogonal to N_1 inverse decays [12, 37] and is not washed out. This wash-out avoidance does not apply to the asymmetry in the τ flavor. Therefore, as we have verified, in all cases we have considered the effect is negligible, since a τ-dominated scenario is always realized.

Let us now calculate the final asymmetry in some interesting cases.

3.1 $\Omega = R_{13}$

The first case we consider is $\Omega = R_{13}$, implying $A_{12} = A_{23} = 0$ in the Eq. (40). As we said already, it is easy to check from the Eq. (26) that $\varepsilon_{2\tau} = 0$ and therefore there is no asymmetry production from N_2-decays even if $M_2 \lesssim 10^{12}\,{\rm GeV}$. On the other hand, one obtains

$$r_{3\tau} = -\frac{2}{3}\,\frac{\sqrt{m_1\,m_3}\,(m_3 - m_1)}{\widetilde{m}_3\,m_{\rm atm}}\,\omega_{31}\,\sqrt{1-\omega_{31}^2}\,c_{12}\,c_{23}^2\,c_{13}\,\Delta\,, \tag{47}$$



Figure 1: Dependence of different quantities on K_1 for $m_1/m_{\rm atm} = 0.1$, $s_{13} = 0.2$, $\delta = -\pi/2$ and real $\Omega = R_{13}$ with $\omega_{31} < 0$. Left panel: projectors $P^0_{1\alpha}$ and $r_{1\alpha}$; central panel: $\xi_{1\alpha}$ and ξ_1 as defined in Eq. (43) for thermal (thin) and vanishing (thick) initial abundance; right panel: lower bound on M_1 for thermal (thin solid) and vanishing (thick solid) initial abundance compared with the one-flavor approximation result (dash-dotted line) obtained for complex $\Omega = R_{13}$. In the squared region the condition Eq. (32) is not satisfied and in the hatched region even the more conservative condition Eq. (33) is not satisfied. The dotted lines (thick for vanishing initial abundance and thin for thermal initial abundance) correspond still to a real $\Omega = R_{13}$ but this time $\delta = 0$ while the only non vanishing low energy phase is the Majorana phase $\Phi_1 = -\pi/2$.

essentially the same expression as for $r_{1\tau}$ but with $\widetilde{m}_1$ replaced by $\widetilde{m}_3$. Therefore, for $M_3 \lesssim 10^{12}\,{\rm GeV}$, one has to worry about a potential non-negligible contribution from N_3 decays. However, when the wash-out from N_1 and N_2 inverse processes is taken into account, see Eq. (46), we always find that the contribution from N_3-decays is negligible and the N_1-dominated scenario holds.

The results are shown in Fig. 1 for $s_{13} = 0.20$, $\delta = -\pi/2$ and $m_1/m_{\rm atm} = 0.1$, a choice of values that approximately maximizes the final asymmetry and yields the lower bound $M_1^{\rm min}(K_1)$. In the left panel we show the tree level projectors $P^0_{1\alpha}$ and the $r_{1\alpha}$'s. It can be seen how for $K_1 \gg 10$ one has $P^0_{1\tau} \simeq P^0_{1,e+\mu} \simeq 1/2$, while for $K_1 \sim 10$ one has $P^0_{1\tau} \ll P^0_{1,e+\mu}$. In the central panel ξ_1 and the $\xi_{1\alpha}$'s are plotted and one can see how for $K_1 \simeq 10$ a τ-dominance is realized. Finally, in the right panel, we show $M_1^{\rm min}(K_1)$ and we compare it with the lower bound in the unflavored approximation obtained for

$\Omega = R_{13}$ (in this case Ω cannot be real) [11]. One can see how, at $K_1 \gg 10$, the asymmetry production rapidly dies, so that $\xi_1 \to 0$ and $M_1^{\rm min}(K_1) \to \infty$. Notice that we plotted the lower bound both for initial thermal N_1-abundance and for initial vanishing N_1-abundance. We also indicated $K_\star$, defined as that value of K_1 such that for $K_1 \gtrsim K_\star$ the dependence on the initial conditions can be neglected and the strong wash-out regime holds. One can notice that the intermediate regime between a fully flavored regime and the unflavored regime, the squared area, is quite extended. In this regime corrections to the results we are showing, obtained in the fully flavored regime, are expected, in a way that the unflavored regime should be recovered for $M_1 \to 10^{12}\,{\rm GeV}$. In this limit the asymmetry production has to switch off and therefore one expects that the lower bound on M_1 has to become more restrictive and eventually, for $M_1 \to 10^{12}\,{\rm GeV}$, the allowed region has to close up. Therefore, one can see that there is no allowed region in the strong wash-out regime. The hatched area, where corrections cannot be excluded within current theoretical uncertainties, cuts away almost completely any allowed region even in the weak wash-out regime. In conclusion, the allowed region where one can safely rely on the fully flavored regime according to current calculations, is very restricted and confined only to a small region in the weak wash-out regime.

3.2 $M_3 \gg 10^{14}\,{\rm GeV}$

The second case we consider is the limit $M_3 \gg 10^{14}\,{\rm GeV}$. In this limit one has necessarily $m_1 \ll m_{\rm sol}$, implying $m_3 \simeq m_{\rm atm}$, and also [49, 50, 51]

$$\Omega = \begin{pmatrix} 0 & 0 & 1 \\ \sqrt{1-\Omega_{31}^2} & -\Omega_{31} & 0 \\ \Omega_{31} & \sqrt{1-\Omega_{31}^2} & 0 \end{pmatrix} . \tag{48}$$

Notice that this particular form of Ω corresponds to set $\omega_{32} = 1$ and $\omega_{21} = 1$ in the Eq. (21). Now in the expression for $r_{1\tau}$ (cf. Eq. (40)) one has $A_{12} = A_{13} = 0$ and therefore

$$r_{1\tau} \simeq \frac{m_{\rm atm}}{\widetilde{m}_1}\sqrt{\frac{m_2}{m_{\rm atm}}}\left(1-\frac{m_2}{m_{\rm atm}}\right)\,\Omega_{31}\,\sqrt{1-\Omega_{31}^2}\,c_{23}^2\,c_{13}\,s_{12}\,\Delta\,. \tag{49}$$

If $M_2 \gtrsim 10^{12}\,{\rm GeV}$, there is no contribution from the next-to-lightest RH neutrino decays anyway, since these occur in the unflavored regime where $\varepsilon_2 \simeq 0$. On the other hand, if $M_2 \lesssim 10^{12}\,{\rm GeV}$, then one has to worry about a (flavored) asymmetry generation from N_2-decays. A calculation of $\varepsilon_{2\alpha}$ shows that the first term in the Eq. (26) vanishes while the second term gives

$$r_{2\tau} = -\frac{2}{3}\,\frac{m_{\rm atm}}{\widetilde{m}_2}\sqrt{\frac{m_2}{m_{\rm atm}}}\left(1-\frac{m_2}{m_{\rm atm}}\right)\,\Omega_{31}\,\sqrt{1-\Omega_{31}^2}\,c_{23}^2\,s_{12}\,\Delta\,. \tag{50}$$



Figure 2: Plots as in Fig. 1 but for the case $M_3 \gg 10^{14}\,\text{GeV}$, corresponding to the special form of Ω in the Eq. (48). Here we are moreover assuming $M_2 \gtrsim 10^{12}\,\text{GeV}$, normal hierarchy. The lower bound $M_1^{\text{min}}(K_1)$ is obtained for $\omega_{31} > 0$ and $\delta = \pi/2$.

This is an example of how the second term in the Eq. (26) is not suppressed in the HL like the first term. However, like for the contribution from N_3-decays in the case $\Omega = R_{13}$, when the wash-out from N_1-inverse processes is taken into account one finds $N^{\rm f}_{B-L}\big|_{N_2} \ll N^{\rm f}_{B-L}\big|_{N_1}$ and a N_1-dominated scenario is realized anyway.

Notice that there is a strong dependence whether one assumes a normal or an inverted hierarchy. For normal hierarchy the results are shown in Fig. 2 for $\omega_{31} > 0$ and $\delta = \pi/2$. For inverted hierarchy the asymmetry is so suppressed that there is no allowed region. This means that for any choice of the parameters one always obtains $M_1^{\text{min}} \gtrsim 10^{12}\,\text{GeV}$.

Notice that results for δ-leptogenesis, in this particular case where $M_3 \gg 10^{14}\,\text{GeV}$, have been recently presented in [23] for vanishing initial N_1 abundance. For example in [23] the authors obtain a lower bound $\sin\theta_{13} \gtrsim 0.09$ imposing the existence of an allowed region for $M_1 \lesssim 5 \times 10^{11}\,\text{GeV}$ while we would obtain $\sin\theta_{13} \gtrsim 0.05$. The difference is probably due to a ($\sim 30\%$) more conservative lower bound that we are using on $\overline{M}_1$ (see Eq. (42)), a difference in the employed value of $m_\star$ (see Eq. (23)), only partly understood in terms of the different convention for the Higgs v.e.v v. There is also a difference in the employed efficiency factor in the strong wash-out regime that, in our case, is about a factor 2 larger. Another likely minor source of difference is that we are not accounting for the effect of spectator processes encoded in the matrix A that relates the $B/3 - L_\alpha$ asymmetries to the L_α asymmetries [12]. However, notice that here we do not want to emphasize too much a precise value of this lower bound on $\sin\theta_{13}$, since we believe this is anyway affected by much larger theoretical uncertainties on the validity of the fully flavored regime. It is however a good way to compare our results with those presented in [23].

3.3 $\Omega = R_{12}$

The third case we consider is $\Omega = R_{12}$. This time one has $A_{13} = A_{23} = 0$ in the Eq.(40). In the case of normal hierarchy the *CP* asymmetry, compared to the case $\Omega = R_{13}$, is suppressed by a factor $(m_{\rm sol}/m_{\rm atm})^{3/2}$, while it is essentially the same for inverted hierarchy. The projectors present very similar features to the case $\Omega = R_{13}$. One can also again calculate, for $M_2 \lesssim 10^{12}\,{\rm GeV}$, the contribution from N_2-decays to the final asymmetry and again one finds that the first term in the Eq. (26) vanishes, while the second produces a term $\propto M_1$, so that

$$r_{2\tau} = \frac{2}{3}\,\frac{\sqrt{m_1\,m_2}\,(m_2 - m_1)}{\widetilde{m}_2\,m_{\rm atm}}\,\omega_{21}\,\sqrt{1-\omega_{21}^2}\,s_{23}\,c_{23}\,\Delta\,. \tag{51}$$

When the efficiency factors are taken into account, one finds that only in the case of normal hierarchy the contribution to the final asymmetry from N_2-decays can be comparable to that one from N_1-decays. However, in this case both productions are suppressed and there is no allowed region anyway in the end. In the case of inverted hierarchy, the contribution from N_2-decays is always negligible compared to that one from N_1-decays. Notice, moreover, that $\varepsilon_{3\alpha} = 0$ and therefore there is no contribution from N_3-decays. In conclusion, for $\Omega = R_{12}$, the lower bound on M_1 for normal hierarchy is much more restrictive than in the case $\Omega = R_{13}$, while it is very similar for inverted hierarchy. A production from the two heavier RH neutrinos can be neglected and the N_1-dominated scenario always holds when the asymmetry is maximized.

3.4 $\Omega = R_{23}$

The last interesting case is $\Omega = R_{23}$. From the Eq. (37) one can easily check that $\varepsilon_{1\alpha} = 0$. One can also easily check that, contrarily to the case $\Omega = R_{12}$, the second term in the Eq. (26) vanishes while the first term does not and yields

$$r'_{2\tau} \equiv \frac{\varepsilon_{2\tau}}{\bar{\varepsilon}(M_2)} = \frac{\sqrt{m_2\,m_3}\,(m_3 - m_2)}{\widetilde{m}_2\,m_{\rm atm}}\,\omega_{32}\,\sqrt{1-\omega_{32}^2}\,s_{12}\,c_{23}^2\,c_{13}\,\Delta\,. \tag{52}$$

Notice that this time $\varepsilon_{2\tau} \propto M_2$ and actually, more generally, one can see that this expression is obtained from the Eq. (40) for $r_{1\tau}$ in the case $\Omega = R_{13}$, just with the replacement



Figure 3: Dependence of different quantities on K_2 for $m_1 = 0$, $s_{13} = 0.2$, $\delta = \pi/2$ and real $\Omega = R_{23}$ with $\omega_{32} > 0$. Left panel: projectors $P^0_{2\alpha}$ and quantities $r'_{2\alpha}$; central panel: $\xi_{2\alpha}$ and ξ_2 for thermal (thin) and vanishing (thick) initial abundance; right panel: lower bound on M_2 for thermal (thin solid) and vanishing (thick solid) abundance compared with the one-flavor approximation result (dash-dotted line) as obtained in [11].

$(M_1, \widetilde{m}_1) \to (M_2, \widetilde{m}_2)$. At the same time one has $K_1 = m_1/m_\star$ so that the wash-out from N_1-inverse processes vanishes for $m_1 \to 0$. For $M_3 \lesssim 10^{12}\,\mathrm{GeV}$ one has to worry about a possible contribution to the asymmetry also from N_3-decays. A straightforward calculation shows that $\varepsilon_{3\alpha} = (2/3)\varepsilon_{2\alpha}$ and therefore an asymmetry is produced at $T \sim M_3$. However, we verified, once more, that the wash-out from N_2-inverse processes is always strong enough that the contribution to the final asymmetry from N_3-decays is negligible.

In complete analogy with the unflavored case [11], one has that the lower bound $M_1^{\rm min}(K_1)$ is replaced by a lower bound $M_2^{\rm min}(K_2)$ obtained for $\omega_{32} > 0$ and shown in the right panel of Fig. 3. One can see that also in this case, within the validity of the condition Eq. (32), the allowed region is constrained to a small portion falling in the weak wash-out regime. Assuming the very conservative condition of validity for the fully flavored regime, outside the squared and hatched regions, there is no allowed region even in the weak wash-out regime.

One can wonder whether there is some choice of Ω, beyond the special cases we analyzed, where the final asymmetry is much higher and the lower bound on M_1 much more relaxed, especially in the strong wash-out regime. We have checked different intermediate cases and we can exclude such a possibility. Therefore, the lower bound shown in Fig. 1

has to be considered, with good approximation, the lowest bound for any choice of real Ω.

Another legitimate doubt is whether, going beyond the approximations we made, the lower bound in Fig. 1 can be considerably relaxed. However, the inclusion of non resonant $\Delta L = 2$ or $\Delta L = 1$ scattering does not produce large corrections. Recently the effect of the off-diagonal terms in the A matrix has been considered, but it has been shown that it does not produce any relevant change in the final asymmetry [18].

Relevant corrections, as already pointed out, can come only from a full quantum kinetic treatment, that should describe accurately the transition between the unflavored regime and the fully flavored regime.

The same kind of considerations holds for the N_2-dominated scenario, realized for $\Omega = R_{23}$. As soon as Ω deviates from R_{23}, the wash-out from N_1 inverse processes comes into play suppressing the final asymmetry and at the same time $\varepsilon_{2\tau}$ gets also suppressed. Therefore, the lower bound on M_2 is necessarily obtained for $\Omega = R_{23}$ in complete analogy with the unflavored approximation [11].

In conclusion δ-leptogenesis in the HL is severely constrained, confirming the conclusions of [9] and [21]. In particular, imposing independence of the initial conditions, then not even a marginal allowed region seems to survive. Notice moreover that all plots have been obtained for $s_{13} = 0.2$, the current $3\,\sigma$ upper bound value. Assuming that for values of M_1 above the condition Eq. (32) the unflavored regime is quickly recovered and therefore that the asymmetry production quickly switches off, then a one-order-of-magnitude improvement of the upper bound on $\sin\theta_{13}$ would essentially completely rule out δ-leptogenesis in the HL, even the marginally allowed regions falling in the weak wash-out regime.

Therefore, in the next section, we will consider the effect of close heavy neutrino masses in enhancing the CP asymmetries and relaxing the lower bounds on M_1, M_2 and the related one on $T_{\rm reh}$. In the end of this section we want to mention that in the more general case of real Ω with non-vanishing Majorana phases, an upper bound $m_1 \lesssim 0.1\,{\rm ev}$ has been obtained in the fully flavored regime [24]. This bound clearly applies also to δ-leptogenesis, but in this case, considering the results we have obtained and the expected quantum kinetic corrections to the fully flavored regime, the issue is actually whether an allowed region exists at all in the HL, even for $m_1 = 0$. Therefore, we do not even try to place an upper bound on m_1 in the HL. In the next section, we will show that actually for δ-leptogenesis an upper bound on m_1 holds even in the resonant limit, where the CP asymmetries are maximally enhanced.

4 The degenerate limit

In this section we show that going beyond the HL the lower bound on M_1 (or on M_2) can be considerably relaxed. Nevertheless, we will see that some interesting constraints on the involved parameters still apply. For simplicity, we can assume a full three-flavor regime holding for M_1 (or M_2) $\ll 10^9\,\mathrm{GeV}$, when also the muon-Yukawa interactions are faster than inverse decays. Therefore now, when we sum over the flavor index, it has to be meant $\alpha = e, \mu, \tau$. This assumption simplifies the calculation, since we do not have to describe a transition between the two and the three-flavor regime and because we can completely neglect the effect, envisaged in [12, 37], for which part of the asymmetry produced from N_2-decays is not touched by N_1-inverse decays. Indeed in a two-flavor regime, even though in the HL we have found that this effect is negligible in all cases we considered because a τ-dominance is always realized, in the DL it can become more relevant because the asymmetry is not necessarily produced dominantly in the τ-flavor.

In order to go beyond the HL, it is convenient to introduce the quantities

$$\delta_{ji} \equiv \frac{M_j - M_i}{M_i} = \sqrt{\frac{x_j}{x_i}} - 1\,. \tag{53}$$

We are interested in the degenerate limit (DL), where at least one δ_{ji} is small enough that both the asymmetry production from decays and the wash-out from inverse processes of the N_i's and of the N_j's can be approximately treated as if they occur at the same temperature, so that they can be simply added up. The DL is a good approximation for $|\delta_{ji}| \lesssim 0.01$ [5]. If $i, j \neq 3$ and $M_3 \gg M_2 \simeq M_1$ then one has a partial DL and in this case the efficiency factors can be approximated, for thermal initial abundance, as [5]

$$\kappa^{\rm f}_{i\alpha} \simeq \kappa^{\rm f}_{j\alpha} \simeq \kappa(K_{i\alpha} + K_{j\alpha})\,. \tag{54}$$

In all considered cases, it will be always verified $K_{i\alpha} + K_{j\alpha} \gg 1$, so that the strong wash-out regime always applies and there is no need to consider the case of initial vanishing abundance. Another possibility is to have a partial DL with $i, j \neq 1$ so that $M_1 \ll M_2 \simeq M_3$. In this case one has to take into account the wash-out from the lightest RH neutrino and therefore

$$\kappa^{\rm f}_{i\alpha} \simeq \kappa^{\rm f}_{j\alpha} \simeq \kappa(K_{i\alpha} + K_{j\alpha})\, e^{-\frac{3\pi}{8} K_{1\alpha}}\,. \tag{55}$$

Finally, in the full DL, one has $M_1 \simeq M_2 \simeq M_3$ and

$$\kappa^{\rm f}_{1\alpha} \simeq \kappa^{\rm f}_{2\alpha} \simeq \kappa^{\rm f}_{3\alpha} \simeq \kappa(K_{1\alpha} + K_{2\alpha} + K_{3\alpha})\,. \tag{56}$$

Let us now calculate the flavored *CP* asymmetries. In the case of real Ω, implying real $(h^\dagger\, h)_{ij} = (h^\dagger h)_{ji}$, the general expression Eq.'s (16) can be conveniently specialized as

$$\varepsilon_{i\alpha} = \frac{3}{16\pi (h^\dagger h)_{ii}} \sum_{j\neq i} (h^\dagger h)_{ij}\, \mathrm{Im}\,[h^\star_{\alpha i}\, h_{\alpha j}] \left[\frac{\xi(x_j/x_i)}{\sqrt{x_j/x_i}} + \frac{2}{3(x_j/x_i - 1)}\right] . \tag{57}$$

In the DL one has approximately $\xi(x_j/x_i) \simeq 1/(3\,\delta_{ji})$ and consequently

$$\varepsilon_{i\alpha} \simeq \frac{1}{8\,\pi (h^\dagger h)_{ii}} \sum_{j\neq i} (h^\dagger h)_{ij}\, \mathrm{Im}\,[h^\star_{\alpha i}\, h_{\alpha j}]\; \delta_{ji}^{-1}. \tag{58}$$

We can again express the neutrino Yukawa coupling matrix through the orthogonal representation. This time the presence of the factor δ_{ji}^{-1} does not allow to remove the sum on j, as it has been possible in the HL in order to derive the Eq. (37). However, considering the same special cases studied in the HL, only one term $j \neq i$ survives and we can write

$$\varepsilon_{i\alpha} \simeq \frac{2\,\bar{\varepsilon}(M_i)}{3\,\delta_{ji}} \sum_{n,h<l} \frac{m_n\,\sqrt{m_h\, m_l}}{\widetilde{m}_i\, m_{\rm atm}}\, \Omega_{ni}\, \Omega_{nj}\, [\Omega_{hi}\, \Omega_{lj} - \Omega_{li}\, \Omega_{hj}]\, \mathrm{Im}[U^\star_{\alpha h}\, U_{\alpha l}]\, . \tag{59}$$

The same expression holds for $\varepsilon_{j\alpha}$ simply exchanging the i and j indexes. We can always choose $j > i$, so that $M_j \geq M_i$. In all the particular cases we will consider it is realized $\varepsilon_{k\alpha} = 0$, for $k \neq i, j$, and moreover the following simplifications apply:

$$\sum_n m_n\, \Omega_{ni}\, \Omega_{nj} = (m_q - m_p)\, \Omega_{ji}\, \Omega_{jj} \quad \text{and} \quad \sum_{h<l} \sqrt{m_h\, m_l}\, [\Omega_{hi}\, \Omega_{lj} - \Omega_{li}\, \Omega_{hj}] = \sqrt{m_q\, m_p}\, , \tag{60}$$

with $q > p$. Except for $M_3 \gg 10^{14}\,\mathrm{GeV}$, in the other cases one has $q = j$ and $p = i$. The final asymmetry can then be expressed as

$$N^{\rm f}_{B-L} \simeq \sum_\alpha (\varepsilon_{i\alpha} + \varepsilon_{j\alpha})\, \kappa^{\rm f}_\alpha(K_{i\alpha} + K_{j\alpha}, K_{k\alpha}) = \frac{\bar{\varepsilon}(M_i)}{3\,\delta_{ji}}\, g(m_1, \Omega_{ji}, \theta_{13}, \delta)\, \Delta\, , \tag{61}$$

where

$$\begin{aligned} g(m_1, \Omega_{ji}, \theta_{13}, \delta) \;\; &\equiv \;\; \frac{2\, K_{\rm atm}\, (K_i + K_j)}{K_i\, K_j}\, \frac{(m_q - m_p)\, \sqrt{m_q\, m_p}}{m^2_{\rm atm}}\, \Omega_{ji}\, \sqrt{1 - \Omega^2_{ji}} \\ &\times \sum_\alpha \kappa^{\rm f}_\alpha(K_{i\alpha} + K_{j\alpha}, K_{k\alpha})\, \frac{\mathrm{Im}[U^\star_{\alpha p}\, U_{\alpha q}]}{\Delta} \end{aligned} \tag{62}$$

and where $\kappa^{\rm f}_\alpha(K_{i\alpha} + K_{j\alpha}, K_{k\alpha}) = \kappa^{\rm f}_{i\alpha} = \kappa^{\rm f}_{j\alpha}$ is given by one of the three expressions Eq. (54), Eq. (55) or Eq. (56), according to the particular case. It is interesting to notice that in the full DL the expression (56) holds and as a consequence of the orthogonality of Ω, one has

$$K_{1\alpha} + K_{2\alpha} + K_{3\alpha} = \sum_k \frac{m_k}{m_\star}\,|U_{\alpha k}|^2\,. \tag{63}$$

In the degenerate limit, since U is unitary, this quantity tends to $m/m_\star$, independently of the flavor, and therefore the sum on the flavors in the Eq. (62) tends to vanish. This will contribute, as we will see, to place a stringent upper bound on the absolute neutrino mass scale in the full DL.

It is also worthwhile to notice that the sign of Δ cannot be predicted from the sign of the observed final asymmetry, since the sign of $g(m_1, \Omega_{ji}, \theta_{13}, \delta)$ depends on the sign of Ω_{ji} that is undetermined. Notice also that $\mathrm{Im}[U^\star_{\alpha h}\,U_{\alpha l}]/\Delta$ does not depend on Δ but nevertheless there is a dependence of $g(m_1, \Omega_{ji}, \theta_{13}, \delta)$ on δ and on θ_{13} coming from the tree level projectors $P^0_{i\alpha}$ in the sum $K_{i\alpha} + K_{j\alpha}$. However, in any case, for $\Delta \rightarrow 0$ one has $g(m_1, \Omega_{ji}, \theta_{13}, \delta)\,\Delta \rightarrow 0$, since the final asymmetry has to vanish when $\sin\theta_{13}$ or $\sin\delta$ vanish.

The function $|g(m_1, \Omega_{ji}, \theta_{13}, \delta)|$ can be maximized over Ω_{ji}. Indeed for $m_1 = 0$, since $\kappa < 1$ and $K_i + K_j \leq K_{\rm atm}$, one has $g(m_1 = 0, K_i, \theta_{13}, \delta) < 4$. Increasing m_1 there is a suppression due to the fact that $K_i \geq m_1/m_\star$ and $g_{\rm max}(m_1, \theta_{13}, \delta)$ decreases monotonically. Therefore, for any m_1, there is a lower bound on M_1 given by

$$M_1 \geq M_1^{\rm min}(m_1, \theta_{13}, \delta) \equiv \frac{3\,\overline{M}_1}{g_{\rm max}(m_1, \theta_{13}, \delta)}\,\frac{\delta_{j1}}{|\Delta|}\,. \tag{64}$$

The *CP* asymmetries, and consequently the final asymmetry, are maximally enhanced in the extreme case of resonant leptogenesis [27, 35] when the heavy neutrino mass degeneracy is comparable to the decay widths. This implies approximately to have $\delta^{\rm res}_{ji} \simeq d\,\bar{\varepsilon}(M_i)/3$ with $d = 1 \div 10$, that would correspond to have $\varepsilon_1 = 1/d$ in the unflavored case with maximal phase. This can be taken as a conservative limit that implies, maximizing over δ, a lower bound

$$\sin\theta_{13} \geq \sin\theta_{13}^{\rm min} = \frac{d\,\eta_B^{\rm CMB}\,N_\gamma^{\rm rec}}{a_{\rm sph}\,\max_\delta[g_{\rm max}(m_1, \theta_{13}^{\rm min}, \delta)\,\sin\delta)]}\,. \tag{65}$$

Notice that, within the validity of perturbation theory, one cannot specify which is the exact value of d, that means the value of δ_{ji} above which the expression for the *CP* asymmetries given in the Eq. (25) are valid [52] and therefore there is an uncertainty in the calculation of the maximum enhancement of the asymmetries in the resonant regime.

Let us now specialize the expressions for the four special cases we have already analyzed in the HL.



Figure 4: Case $M_3 \gg 10^{14}\,\mathrm{GeV}$ for normal hierarchy in the DL. Plot of the function $|g(K_1,\theta_{13},\delta)|$ in the limit $\Delta \to 0$. The maximum gives the lower bound on M_1 (see Eq. (67)) and on $\sin\theta_{13}$ (see Eq. (69)).

4.1 $M_3 \gg 10^{14}\,\mathrm{GeV}$

Remember that in this case one has $(h^\dagger h)_{3j} = 0$ implying $\varepsilon_{3\alpha} = 0$, a consequence of the fact that the heaviest RH neutrino decouples. Moreover $m_1 \ll m_{\rm sol}$, such that terms $\propto m_1$ can be neglected, $m_3 \simeq m_{\rm atm}$ and $m_2 \simeq m_{\rm sol}$ for normal hierarchy or $m_2 \simeq m_{\rm atm}\sqrt{1 - m_{\rm sol}^2/m_{\rm atm}^2}$ for inverted hierarchy. Therefore, there is actually no dependence on m_1 in $g(m_1, \Omega_{ji}, \theta_{13}, \delta)$ that we can indicate simply with $g(\Omega_{ji}, \theta_{13}, \delta)$ and that is given by the expression (62) with $(i,j) = (1,2)$ and $(p,q) = (2,3)$ or explicitly

$$\begin{aligned} g(\Omega_{21},\theta_{13},\delta) \;\; &\simeq \;\; \frac{2\,(K_1+K_2)\,K_{\rm atm}}{K_1\,K_2}\,\left(1-\frac{m_2}{m_{\rm atm}}\right)\,\sqrt{\frac{m_2}{m_{\rm atm}}}\,\Omega_{21}\,\sqrt{1-\Omega_{21}^2} \\ &\times \sum_\alpha \kappa(K_{1\alpha}+K_{2\alpha})\,\frac{{\rm Im}[U_{\alpha 2}^\star\,U_{\alpha 3}]}{\Delta}. \end{aligned} \tag{66}$$

In the case of normal hierarchy $|g(\Omega_{21},\theta_{13},\delta)|$ slightly decreases when Δ increases and so the maximum is found for $\Delta = 0$ and in this case the dependence on θ_{13} and on δ disappears. Replacing Ω_{21} with K_1, in Fig. 4 we have plotted $|g(K_1, \Delta = 0)|$ for central values of $m_{\rm sol}$ and $m_{\rm atm}$. Including the errors, one finds $g_{\rm max} \simeq 0.160 \pm 0.005$.

The (3σ) lower bounds on M_1 for normal hierarchy, from the general expression (64), is then given by

$$M_1 \geq 0.9 \times\, 10^{10}\,\mathrm{GeV}\,\frac{\delta_{21}}{|\Delta|}\,. \tag{67}$$

In the case of inverted hierarchy the situation is somehow opposite, since for $\theta_{13} = 0$ the electron flavor contribution vanishes in the Eq. (66) and there is an exact cancellation between the τ and μ contributions. Consequently, the asymmetry increases for increasing values of θ_{13} and thus the maximum is found for $\sin\theta_{13} = 0.2$ while $\delta \simeq \pi/4$. In this case one has that $\max_{\theta_{13},\delta}[g_{\max}(m_1 = 0, \theta_{13}, \delta)\,\Delta] \simeq (9 \pm 2) \times 10^{-8}$, that plugged in the Eq. (64) gives at 3σ

$$M_1 \geq 6 \times 10^{15}\,\mathrm{GeV}\,\delta_{21}\,. \tag{68}$$

It should be remembered that these conditions have been obtained in the three-flavor regime and in the DL and therefore are valid for $M_1 \lesssim 10^9\,\mathrm{GeV}$. This implies $\delta_{21} \lesssim 10^{-1}\,|\Delta|$ for normal hierarchy and $\delta_{21} \lesssim 10^{-7}$ for inverted hierarchy.

Analogously the general expression (65) gives, for normal and inverted hierarchy respectively, the following (3σ) lower bounds on $\sin\theta_{13}$:

$$\sin\theta_{13} \gtrsim 3.3 \times 10^{-7}\,d \quad \text{and} \quad \sin\theta_{13} \gtrsim 0.06\,d. \tag{69}$$

4.2 $\Omega = R_{13}$

In this particular case, the next-to-lightest RH neutrino is decoupled from the other two and this implies that $\varepsilon_{2\alpha} = 0$ for any α and that the $\varepsilon_{1\alpha}$'s do not depend on M_2, in particular they do not get enhanced if $\delta_{21} \to 0$. Therefore, one has necessarily to consider $\delta_{31} \lesssim 0.01$, implying a full DL with all three degenerate RH neutrino masses. The function $g(m_1, \Omega_{ji}, \theta_{13}, \delta)$ is now obtained from the general expression (62) for $j = q = 3$ and $i = p = 1$, or explicitly

$$\begin{aligned} g(m_1, \Omega_{31}, \theta_{13}, \delta) \;\equiv\; & \frac{2\,K_{\mathrm{atm}}\,(K_1 + K_3)}{K_1\,K_3}\,\frac{(m_3 - m_1)\,\sqrt{m_3\,m_1}}{m^2_{\mathrm{atm}}}\,\Omega_{31}\,\sqrt{1 - \Omega^2_{31}} \\ & \times \sum_\alpha \kappa(K_{1\alpha} + K_{2\alpha} + K_{3\alpha})\,\frac{\mathrm{Im}[U^\star_{\alpha 1}\,U_{\alpha 3}]}{\Delta}\,. \end{aligned} \tag{70}$$

It is interesting to notice that in this case an e-dominance is realized. Moreover, one has that the dependence of $|g(m_1, \Omega_{31}, \theta_{13}, \delta)|$ on θ_{13} and δ is slight and the maximum is again for $\Delta = 0$ and for $m_1 = 0$ and one finds $g_{\max}(0) = 0.24 \pm 0.01$ for normal hierarchy and $g_{\max}(0) = (3.1 \pm 0.2) \times 10^{-3}$ for inverted hierarchy, so that the general expression (64) for the lower bound on M_1 gives, at 3σ for normal and inverted hierarchy,

$$M_1 \gtrsim 5.5 \times 10^9\,\mathrm{GeV}\,\frac{\delta_{31}}{|\Delta|} \quad \text{and} \quad M_1 \gtrsim 5 \times 10^{11}\,\mathrm{GeV}\,\frac{\delta_{31}}{|\Delta|}\,, \tag{71}$$



Figure 5: Case $\Omega = R_{13}$ in the full DL. Lower bound on $\sin\theta_{13}$ versus m_1 obtained in resonant leptogenesis for $d = 1$ (solid line) and $d = 10$ (short-dashed line). Values $\sin\theta_{13} > 0.20$ are excluded at $3\,\sigma$ by current experimental data.

while the general expression (65) in resonant leptogenesis gives

$$\sin\theta_{13} \gtrsim 2.3\times10^{-7}\,d \quad \text{and} \quad \sin\theta_{13} \gtrsim 1.5\times10^{-5}\,d\,. \tag{72}$$

Increasing m_1, the value of $g_{\max}(m_1)$ decreases and the lower bound on $\sin\theta_{13}$ in resonant leptogenesis becomes more and more restrictive. This dependence is shown in Fig. 5 both for normal (left panel) and inverted (right panel) hierarchy and for $d = 1$ (solid line) and $d = 10$ (short-dashed line). Very interestingly, imposing the experimental (3σ) upper bound $\sin\theta_{13} \lesssim 0.20$, one obtain the upper bound $m_1 \lesssim (0.2-0.4)\,\mathrm{eV}$, depending on the value of d. This upper bound will become more stringent if the experimental upper bound on $\sin\theta_{13}$ will improve, as expected in future experiments in the case of no discovery. The most stringent experimental upper bound that can be hopefully reached in future with neutrino factories is approximately $\sin\theta_{13} < 10^{-3}$ [53]. This asymptotical upper bound is also shown in Fig. 5 and would imply an upper bound $m_1 \lesssim (0.05-0.1)\,\mathrm{eV}$ for normal hierarchy and $m_1 \lesssim (0.03-0.08)\,\mathrm{eV}$ for inverted hierarchy. Therefore, an interesting interplay between two measurable quantities is realized and this makes δ-leptogenesis falsifiable independently of the RH neutrino mass spectrum.

In the more conservative case of normal hierarchy, see left panel of Fig. 5, a good



Figure 6: Case $\Omega = R_{12}$ in the partial DL. Lower bound on $\sin\theta_{13}$ versus m_1 obtained in resonant leptogenesis. Same conventions as in the previous figure.

approximation is given by the fit

$$m_1 \lesssim 0.6\,\left(\sin\theta_{13} - 2.3\times 10^{-7}\right)^{0.25}\,{\rm eV}\,. \tag{73}$$

It is interesting that this upper bound holds in the extreme case of resonant leptogenesis and therefore holds for any RH neutrino spectrum. However, we have to verify whether it holds also for a different choice of Ω.

4.3 $\Omega = R_{12}$

The situation for $\Omega = R_{12}$ is quite different compared to the previous cases. Now one has $i = p = 1$ and $j = q = 2$ and it is possible to have both a partial DL with $10^{14}\,{\rm GeV} \gtrsim M_3 \gg M_2 \simeq M_1$ and a full DL. In the first case, the general expression Eq. (62) becomes

$$\begin{aligned} g(m_1,\Omega_{21},\theta_{13},\delta) \;\; &\equiv \;\; \frac{2\,K_{\rm atm}\,(K_1+K_2)}{K_1\,K_2}\,\frac{(m_2-m_1)\,\sqrt{m_2\,m_1}}{m_{\rm atm}^2}\,\Omega_{21}\,\sqrt{1-\Omega_{21}^2} \\ &\times \sum_\alpha \kappa(K_{1\alpha}+K_{2\alpha})\,\frac{{\rm Im}[U_{\alpha 1}^\star\,U_{\alpha 2}]}{\Delta}\,. \end{aligned} \tag{74}$$

This time the contribution from the electron flavor vanishes. Furthermore, for normal hierarchy, there is an almost perfect cancellation between the μ and the τ contribution. In the left panel of Fig. 6 we show the lower bound on $\sin\theta_{13}$ versus m_1 and one can



Figure 7: Case $\Omega = R_{12}$ in the full DL. Lower bound on $\sin\theta_{13}$ versus m_1 obtained in resonant leptogenesis. Same conventions as in the previous figures.

see how, compared to the previous case $\Omega = R_{13}$, this is much more restrictive. In particular, imposing $\sin\theta_{13} < 0.2$, one obtains now a much more stringent upper bound $m_1 \lesssim 0.06\,\mathrm{eV}$. On the other hand, for inverted hierarchy, the cancellation between the μ and the τ flavor does not occur and one has a lower bound on $\sin\theta_{13}$, for $m_1 \ll 0.01\,\mathrm{eV}$, shown in the right panel of Fig. 6, that is very similar to what has been obtained in the case $\Omega = R_{13}$. However, now there is no flavor cancellation for increasing values of m_1, because $K_{1\alpha} + K_{2\alpha}$ does not tend to a common value like $\sum_j K_{j\alpha}$. Therefore, one can see in Fig. 6 that this time the upper bound on m_1 is much looser, both compared to normal hierarchy and compared to $\Omega = R_{13}$.

In the full DL, the flavor cancellation at large m_1 occurs and the results are shown in Fig. 7. One can see how now for normal hierarchy the upper bound on m_1 is even much more restrictive and, for inverted hierarchy, one has a situation that is similar to the case $\Omega = R_{13}$.

4.4 $\Omega = R_{23}$

In this case the lightest RH neutrino decouples and $\varepsilon_{1\alpha} = 0$, independently of M_1. Therefore, there is no contribution to the final asymmetry from N_1 decays. On the other hand $\varepsilon_{2\alpha}$ and $\varepsilon_{3\alpha}$ do not vanish and therefore there is a contribution from the decays of the two



Figure 8: Case $\Omega = R_{23}$ in the full DL. Lower bound on $\sin\theta_{13}$ versus m_1 obtained in resonant leptogenesis. Same conventions as in the previous figures.

heavier RH neutrinos. Still N_1 inverse processes have to be taken into account since they contribute to the wash-out. There are two different possibilities.

In a full DL the wash-out from N_1 inverse decays just cumulates with the wash-out from the two heavier. Therefore, this time, in the expression Eq. (62), one has $i = p = 2$ and $j = q = 3$ and $\kappa_\alpha^{\rm f} = \kappa(K_{1\alpha} + K_{2\alpha} + K_{3\alpha})$, explicitly

$$\begin{aligned} g(m_1, \Omega_{32}, \sin\theta_{13}, \sin\delta) \;\; &\equiv \;\; \frac{2\,K_{\rm atm}\,(K_2 + K_3)}{K_2\,K_3}\,\frac{(m_3 - m_2)\,\sqrt{m_3\,m_2}}{m_{\rm atm}^2}\,\Omega_{32}\,\sqrt{1 - \Omega_{32}^2} \\ &\times \sum_\alpha \kappa(K_{1\alpha} + K_{2\alpha} + K_{3\alpha})\,\frac{{\rm Im}[U_{\alpha 2}^\star\,U_{\alpha 3}]}{\Delta}\,. \end{aligned} \tag{75}$$

In Fig. 8 we show the dependence of the $\sin\theta_{13}$ lower bound on m_1. This time there is a bigger suppression than in the case $\Omega = R_{13}$, both for normal and for inverted hierarchy.

In the case $M_1 \ll M_2 \simeq M_3$ one has

$$\begin{aligned} g(m_1, \Omega_{32}, \sin\theta_{13}, \sin\delta) \;\; &\equiv \;\; \frac{2\,K_{\rm atm}\,(K_2 + K_3)}{K_2\,K_3}\,\frac{(m_3 - m_2)\,\sqrt{m_3\,m_2}}{m_{\rm atm}^2}\,\Omega_{32}\,\sqrt{1 - \Omega_{32}^2} \\ &\times \sum_\alpha \kappa(K_{2\alpha} + K_{3\alpha})\,e^{-\frac{3\pi}{8}\,K_{1\alpha}}\,\frac{{\rm Im}[U_{\alpha 2}^\star\,U_{\alpha 3}]}{\Delta}\,. \end{aligned} \tag{76}$$

The dependence of the lower bound on $\sin\theta_{13}$ on m_1 is shown in Fig. 9 for normal hierarchy. In this case the upper bound on m_1 is now slightly less stringent than in the



Figure 9: Case $\Omega = R_{23}$ in the partial DL. Lower bound on $\sin\theta_{13}$ versus m_1 obtained in resonant leptogenesis. Same conventions as in the previous figures.

previous cases. For inverted hierarchy the final asymmetry production is so suppressed that there is no allowed region.

We can conclude this section noticing that these results show that δ-leptogenesis can be falsified. In the case of normal hierarchy, the current upper bound $\sin\theta_{13} \lesssim 0.2$ implies $m_1 \lesssim 0.1\,\mathrm{eV}$, while, in future, a potential upper bound $\sin\theta_{13} \lesssim 10^{-3}$ would imply $m_1 \lesssim \mathcal{O}(0.01\,\mathrm{eV})$, with a more precise determination depending on the possibility of improving the current estimation of the parameter d in resonant leptogenesis.

5 Lights and shadows of δ-leptogenesis

The most attractive feature of δ-leptogenesis is that a non-vanishing Dirac phase, the only see-saw phase that we can realistically hope to discover in future, acts as the only source of *CP* violation responsible for the matter-antimatter asymmetry of the Universe. We think that this feature, despite of the objections that we are going to discuss, provides a strong motivation for δ-leptogenesis.

As we have seen, successful δ-leptogenesis implies stringent conditions on the RH neutrino masses, something quite interesting since they escape conventional experimental information. In particular we have seen that, except for a marginal allowed region in the weak wash-out regime, the HL is non-viable. We also observed that a definite conclusion on the existence of such a marginal allowed region, requires a full quantum kinetic treatment but in any case corrections are expected to shrink this already quite restricted allowed region.

Therefore, δ-leptogenesis motivates models with degenerate RH neutrino masses, with the most extreme limit represented by resonant leptogenesis. Even in this extreme limit however, imposing successful δ-leptogenesis, interesting conditions follow on quantities accessible in low-energy neutrino experiment: $\sin\theta_{13}$, the absolute neutrino mass scale, normal or inverted scheme, the Dirac phase itself. Therefore, an interesting aspect of δ-leptogenesis is that it is falsifiable independently of the heavy neutrino mass spectrum.

There are some objections to δ-leptogenesis. There is no clear theoretical motivation for δ-leptogenesis, more generally to choose a real orthogonal Ω matrix. Apparently, sequential dominated models [54] could represent an interesting theoretical framework. Indeed in [11] it was shown that these models correspond to have an Ω matrix that slightly deviates from the unit matrix or from all the other five that can be obtained from the unit matrix exchanging rows or columns. However, it has been noticed [4, 11] that in the limit $\mathrm{Im}[\Omega] \rightarrow 0$ total CP asymmetries ε_i do not necessarily vanish. Therefore, in this limit and taking vanishing Majorana phases, one does not necessarily obtain δ-leptogenesis. Writing $\Omega^2_{ij} = |\Omega^2_{ij}| \exp[i\,\varphi_{ij}]$, the correct condition to enforce $\varepsilon_i \rightarrow 0$ is to take the limit $\varphi_{ij} \rightarrow 0$. This is a more demanding limit than $\mathrm{Im}[\Omega] \rightarrow 0$ and it is currently not motivated by generic sequential dominated models. This limit is not motivated either by radiative leptogenesis [29] within the context of the minimal flavor violation principle [30], as recently considered in [55, 28]. Therefore, there is no theoretical justification for δ-leptogenesis at the moment.

Another possible objection to δ-leptogenesis is that it cannot be distinguished from the general scenario, where all phases are present, even if a non-vanishing Dirac phase is discovered. Indeed a Dirac phase would give in this case a subdominant contribution. This objection is however related also to the first one. Indeed, since a theoretical model motivating δ-leptogenesis is required anyway, one can hope to find some specific prediction that makes the model testable and δ-leptogenesis together with it. Dirac phase leptogenesis would then become distinguishable from the general scenario, though in an indirect way.

This last objection can be also considered within a more particular case where Ω is still real but Majorana phases are present together with the Dirac phase. It has been noticed that the contribution to the final asymmetry from Majorana phases is in general dominant compared to that one coming from the Dirac phase [9]. In the right panel of Fig. 1 we have compared the result on the M_1 lower bound for $\Omega = R_{13}$ obtained in

δ-leptogenesis with the result when $\text{Im}[\Omega_{ij}] = \delta = 0$ but $\Phi_1 = -\pi/2$ (dotted lines). One can see that in the second case the lower bound is $\sim 2 \div 3$ times more relaxed. This result can be easily understood analytically [22] and actually it can be also observed that there can be exact cancellations between the contribution to the final asymmetry from the Majorana phases and from the Dirac phase.

The presence of cancellations can be somehow regarded as a limit to δ-leptogenesis main motivation, since even though a Dirac phase will be discovered, it is not guaranteed that the observed asymmetry can be explained. This objection is however quite weak since it would be quite strange if Nature disposed a sufficient source of *CP* violation but set up a second source that exactly cancels with the first one while the observed asymmetry is, in the end, explained still by a third one, for example the phases in Ω. On the other hand, we can say that it would be certainly positive for δ-leptogenesis if in future experimental upper bounds on the Majorana phases are placed, for example from $\beta\beta 0\nu$ decay, thus constraining the contribution to the final asymmetry from Majorana phases [22]. This can be also regarded as a further prediction coming from δ-leptogenesis.

In conclusion, we have studied in detail a specific scenario of leptogenesis that is interesting especially in view of the many next planned experiments aiming at a discovery of *CP* violation in neutrino mixing. Despite some important remarks and objections, we think that δ-leptogenesis realizes a very interesting link between a long-standing cosmological puzzle and *CP* violation in neutrino oscillations, one of the most relevant experimental topics in high-energy physics during next years.

Acknowledgments

It is a pleasure to thank S. Petcov for discussions during the Neutrino Oscillation Workshop 2006, held in Conca della Specchiulla (Italy). We also wish to thank G. Raffelt for interesting discussions and comments. This work was supported, in part, under the Marie Curie project "Leptogenesis, Seesaw and GUTs," contract No. MEIF-CT-2006-022950.

References

[1] M. Fukugita, T. Yanagida, Phys. Lett. **B 174** (1986) 45.

[2] P. Minkowski, Phys. Lett. B **67** (1977) 421; T. Yanagida, in *Workshop on Unified Theories*, KEK report 79-18 (1979) p. 95; M. Gell-Mann, P. Ramond, R. Slansky, in *Supergravity* (North Holland, Amsterdam, 1979) eds. P. van Nieuwenhuizen, D. Freedman, p. 315; S.L. Glashow, in *1979 Cargese Summer Institute on Quarks and Leptons* (Plenum Press, New York, 1980) p. 687; R. Barbieri, D. V. Nanopoulos, G. Morchio

and F. Strocchi, Phys. Lett. B **90** (1980) 91; R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44** (1980) 912.

[3] V. A. Kuzmin, V. A. Rubakov, M. E. Shaposhnikov, Phys. Lett. **B 155** (1985) 36.

[4] W. Buchmüller, P. Di Bari and M. Plümacher, Nucl. Phys. B **665** (2003) 445.

[5] S. Blanchet and P. Di Bari, JCAP **0606** (2006) 023.

[6] S. Davidson, A. Ibarra, Phys. Lett. **B 535** (2002) 25.

[7] W. Buchmüller, P. Di Bari and M. Plümacher, Nucl. Phys. B **643** (2002) 367.

[8] W. Buchmüller, P. Di Bari and M. Plümacher, Annals Phys. **315** (2005) 305.

[9] S. Blanchet and P. Di Bari, JCAP **03** (2007) 018.

[10] G. F. Giudice, A. Notari, M. Raidal, A. Riotto and A. Strumia, Nucl. Phys. B **685** (2004) 89.

[11] P. Di Bari, Nucl. Phys. B **727** (2005) 318.

[12] R. Barbieri, P. Creminelli, A. Strumia and N. Tetradis, Nucl. Phys. B **575** (2000) 61.

[13] T. Endoh, T. Morozumi and Z. h. Xiong, Prog. Theor. Phys. **111** (2004) 123.

[14] A. Pilaftsis, Phys. Rev. Lett. **95** (2005) 081602; A. Pilaftsis and T. E. J. Underwood, Phys. Rev. D **72** (2005) 113001.

[15] E. Nardi, Y. Nir, E. Roulet and J. Racker, JHEP **0601** (2006) 164.

[16] A. Abada, S. Davidson, F. X. Josse-Michaux, M. Losada and A. Riotto, JCAP **0604** (2006) 004.

[17] A. Abada, S. Davidson, A. Ibarra, F. X. Josse-Michaux, M. Losada and A. Riotto, JHEP **0609** (2006) 010.

[18] F. X. Josse-Michaux and A. Abada, arXiv:hep-ph/0703084.

[19] M. Y. Khlopov and A. D. Linde, Phys. Lett. B **138** (1984) 265. J. R. Ellis, J. E. Kim and D. V. Nanopoulos, Phys. Lett. B **145** (1984) 181; K. Kohri, T. Moroi and A. Yotsuyanagi, Phys. Rev. D **73** (2006) 123511.

[20] J. A. Casas and A. Ibarra, Nucl. Phys. B **618** (2001) 171.

[21] S. Antusch and A. M. Teixeira, JCAP **0702** (2007) 024.

[22] S. Pascoli, S. T. Petcov and A. Riotto, Phys. Rev. D **75** (2007) 083511.

[23] S. Pascoli, S. T. Petcov and A. Riotto, arXiv:hep-ph/0611338.

[24] G. C. Branco, R. Gonzalez Felipe and F. R. Joaquim, Phys. Lett. B **645** (2007) 432.

[25] S. Blanchet, P. Di Bari and G. G. Raffelt, JCAP **03** (2007) 012.

[26] L. Covi, E. Roulet, F. Vissani, Phys. Lett. **B 384** (1996) 169.

[27] A. Pilaftsis, Int. J. Mod. Phys. A **14** (1999) 1811.

[28] S. Uhlig, arXiv:hep-ph/0612262.

[29] R. Gonzalez Felipe, F. R. Joaquim and B. M. Nobre, Phys. Rev. D **70** (2004) 085009; G. C. Branco, R. Gonzalez Felipe, F. R. Joaquim and B. M. Nobre, Phys. Lett. B **633** (2006) 336.

[30] V. Cirigliano, B. Grinstein, G. Isidori and M. B. Wise, Nucl. Phys. B **728** (2005) 121.

[31] M. C. Gonzalez-Garcia and M. Maltoni, arXiv:0704.1800 [hep-ph].

[32] B. A. Campbell, S. Davidson, J. R. Ellis and K. A. Olive, Phys. Lett. B **297** (1992) 118; J. M. Cline, K. Kainulainen and K. A. Olive, Phys. Rev. D **49** (1994) 6394.

[33] M. A. Luty, Phys. Rev. D **45** (1992) 455.

[34] M. Plümacher, Z. Phys. **C 74** (1997) 549.

[35] A. Pilaftsis and T. E. J. Underwood, Nucl. Phys. B **692** (2004) 303.

[36] W. Buchmüller and M. Plümacher, Phys. Lett. B **511** (2001) 74.

[37] E. Nardi, Y. Nir, J. Racker and E. Roulet, JHEP **0601** (2006) 068.

[38] D. N. Spergel *et al.* [WMAP Collaboration], arXiv:astro-ph/0603449.

[39] K. S. Babu, C. N. Leung and J. T. Pantaleone, Phys. Lett. B **319** (1993) 191; S. Antusch, J. Kersten, M. Lindner and M. Ratz, Nucl. Phys. B **674** (2003) 401.

[40] S. Eidelman et al., Phys. Lett. **B592**, 1 (2004) (URL:http://pdg.lbl.gov/).

[41] M. Fujii, K. Hamaguchi and T. Yanagida, Phys. Rev. D **65** (2002) 115012.

[42] P. Di Bari, AIP Conf. Proc. **655** (2003) 208 [arXiv:hep-ph/0211175]. P. Di Bari, [arXiv:hep-ph/0406115].

[43] W. Buchmüller, P. Di Bari and M. Plümacher, Phys. Lett. B **547** (2002) 128.

[44] A. De Simone and A. Riotto, JCAP **0702** (2007) 005.

[45] O. Vives, Phys. Rev. D **73** (2006) 073006.

[46] G. Engelhard, Y. Grossman, E. Nardi and Y. Nir, arXiv:hep-ph/0612187.

[47] T. Shindou and T. Yamashita, arXiv:hep-ph/0703183.

[48] J. F. Nieves and P. B. Pal, Phys. Rev. D **36** (1987) 315.

[49] P. H. Frampton, S. L. Glashow and T. Yanagida, Phys. Lett. B **548** (2002) 119.

[50] A. Ibarra and G. G. Ross, Phys. Lett. B **575** (2003) 279.

[51] P. H. Chankowski and K. Turzynski, Phys. Lett. B **570** (2003) 198.

[52] A. Anisimov, A. Broncano and M. Plumacher, Nucl. Phys. B **737** (2006) 176.

[53] P. Huber, M. Lindner and W. Winter, Nucl. Phys. B **645** (2002) 3.

[54] See S. F. King, Rept. Prog. Phys. **67** (2004) 107 and references therein.

[55] V. Cirigliano, G. Isidori and V. Porretti, Nucl. Phys. B **763** (2007) 228. G. C. Branco, A. J. Buras, S. Jager, S. Uhlig and A. Weiler, arXiv:hep-ph/0609067.